arXiv:1205.0172v1 [math.PR] 1 May 2012

Bifurcations of stochastic differential equations with singular diffusion coefficients

Jonathan Touboul*[a,b], Gilles Wainrib[c]

[a]*The Mathematical Neuroscience Laboratory, CIRB*[1]*, Collège de France, Paris, France*
[b]*BANG Laboratory, INRIA Paris Rocquencourt.*
[c]*LAGA, Paris 13 University, Villetaneuse, France*

Abstract

In this article, we address the dynamics and bifurcations of a wide class of stochastic differential equations around singular points where both the drift and diffusion functions vanish. We apply these results to stochastic versions of the pitchfork, Hopf and saddle-node bifurcations. According to the Hölder coefficient of the diffusion function around the singular point, we identify different regimes, and the multiplicative noise case appears as a singular transition.

Keywords: bifurcations, noise, universal unfolding, stochastic stability, (quasi)-stationary distributions, normal forms

---

1. Introduction

A large class of phenomena can be described using non-linear ordinary differential equations (ODEs), in such distinct domains as economics, physics or biology. Many analytical methods have been developed to characterize the solutions of such systems. In particular, the widely applied theory of bifurcations allows a qualitative characterization of the asymptotic regimes of nonlinear dynamical systems. This theory ensures that the number or stability of fixed points and periodic orbit can change according to a few generic scenarii, and the system at the bifurcation point can be locally mapped to a simple polynomial dynamical system, called normal form [1, 2]. This theory is very handy to study nonlinear differential equations and is almost systematically used to characterize the asymptotic behavior of complicated systems. Investigating the impact of noisy perturbations on such systems is hence of great interest, and is currently an active field of research. In the last decades, it has been shown in many different areas that applying noise on a non-linear system can lead to many counter-intuitive phenomena, such as noise-induced stabilization [3], oscillations [4] or transport [5, 6] and stochastic resonance [7, 8]. From a mathematical perspective, understanding the interplays between non-linearities and noisy perturbations is a great challenge, with many applications. Several tools have been introduced, ranging from the theory of random dynamical systems (RDS) [9, 10], the study of moment equations [11, 12] or multiscale stochastic methods for slow-fast systems [13, 14]. Unfortunately no systematic method in the flavor of bifurcation theory for the analysis of the dynamics of nonlinear ODEs exist, and this is the central problematic of the present manuscript.

---

*Corresponding Author: `jonathan.touboul@college-de-france.fr`

[1] CNRS UMR 7241, INSERM U1050, UPMC ED 158. MEMOLIFE & PSL*

We focus here on the dynamics of stochastic differential equations (SDE) in which both the drift and the diffusion functions vanish at some particular point, called singular point. This type of equations may appear for example in diffusion approximations of population models [15] or in the modeling of noisy neural conductances [16]. The question we address is how the interplay between the drift and the shape of the diffusion function affects the behavior of the system. An important contribution of RDS theory to the field of stochastic bifurcations is to distinguish between phenomenological (P) bifurcations, characterized by a change in the shape of the invariant measure, and dynamical (D) bifurcations, characterized by a sign change of the stochastic Lyapunov exponent. We investigate the questions of stochastic stability and bifurcations (P and D) in a comprehensive manner, using combinations of appropriate tools such as Lyapunov functionals and exponents, stationary solutions of Fokker-Planck equations and Feller scale functions (see e.g. [17] for an extensive account on these concepts). Mathematically, the choice of a vanishing diffusion coefficient provides a framework to study subtle competitions between the drift and the noise, leading to a rich and generic phenomenology.

With the aim of characterizing the behavior of the solutions of SDEs with drift given by the normal form of usual bifurcations, we will start by analyzing a general one dimensional stochastic differential equation

$$dx_t = f(x_t)\,dt + \gamma(x_t)\,dW_t$$

where $W = (W_t)_{t\geq 0}$ is a Brownian motion defined on a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$ endowed with the natural filtration $(\mathcal{F}_t)_t$ of W. Notice that several classical normal forms can be reformulated in this setting and our general results will be applied to the cases of the pitchfork, Hopf or saddle-node bifurcations in section 3. We assume that there exists a singular point x^* such that $f(x^*) = \gamma(x^*) = 0$. The system has a trivial solution distributed as a Dirac measure localized at the singular point x^*, and we shall address the stochastic stability of these solutions. In contrast to the case of ODEs, there are at least three different notions of stochastic stability:

1. The *almost sure exponential stability* defined by the property

   $$\limsup_{t\to\infty} \frac{1}{t}\log|x_t - x^*| < 0 \quad a.s.$$

   for any initial condition $X_0 \in \mathbb{R}$. The *almost sure exponentially instability* defined by the property

   $$\liminf_{t\to\infty} \frac{1}{t}\log|x_t - x^*| > 0 \quad a.s.$$

   for any initial condition $X_0 \in \mathbb{R}$ (see e.g. [18]). This definition implies a strong notion of stability close to the usual exponential stability property of deterministic dynamical systems: almost all trajectories exponentially fast converge to or diverge from x^*. In our stochastic context, this notion can appear quite restrictive, and we will hence also discuss a weaker notion of stability:
2. The *stochastic stability* (or stability in probability), defined by the property that for all $\mu \in (0, 1)$ and $r > 0$ there exists δ depending on μ and r such that

   $$\mathbb{P}(\{|x_t - x^*| < r \ \ \forall t \geq 0\}) \geq 1 - \mu$$

   whenever $|x_0 - x^*| < \delta$ and otherwise it is said to be unstable in probability. The solution x^* is *stochastically asymptotically stable* (or asymptotically stable in probability) if it is stable in probability and for every $\mu \in (0,1)$ there exists η_0 depending on μ such that

$$\mathbb{P}(\lim_{t\to\infty} |x_t - x^*| = 0) \geq 1 - \mu$$

whenever $|x_0 - x^*| < \eta_0$.

3. The *stability (resp. instability) in the first approximation* of a stationary solution $\bar{X}$ [17, Chap. 7.1], notion widely used in the RDS theory [9], consists in that the identically zero solution of the linearized equation around $\bar{X}$ is almost surely exponentially stable (resp. unstable). This criterion will be chiefly used when dealing with non-Dirac stationary solutions.

To be more specific, we now consider that $x^* = 0$ and assume that $f(x)$ is a smooth function, which is often the case in applications and a basis of the deterministic bifurcation theory. In detail, we assume that $f(x) = \lambda\, x + g(x)$ where $g(x) = o(x)$ at $x = 0$ (i.e. $f'(0) = \lambda$), and hence consider the following class of SDEs on $\mathbb{R}$:

$$dx_t = (\lambda x_t + g(x_t))\, dt + \gamma(x_t) dW_t \tag{1}$$

with $\gamma(x) \sim \sigma|x|^\alpha$ at $x = 0$. The behavior of the solutions will be described as a function of the local behavior of g and γ around 0 similarly to the deterministic case, and these properties will precisely characterize the stability of the solutions. However, in contrast with the deterministic case, the existence of stationary distributions will depend on the properties of g and γ at infinity. Intuitively, this is due to the fact that stationary solutions can reach values far from the singular point, where the local behavior of the solution is no more relevant. In Figure 1, the main question of the paper is schematically described, in the setting of a particle in a double well potential subjected to noisy fluctuations that vanish on the saddle point and increase as a function of the distance. According to the shape of the noise function γ (mainly the parameter α), the particle will have two distinct fates : either it falls into the well and keeps fluctuating without ever reaching the saddle point, or it eventually reaches it after some finite time and remains steady for all time.

The paper is organized as follows. We start by analyzing in detail in section 2 the behavior of the solution of a general diffusion equation (1). This section contains several abstract results summarized in Theorem 1 and a general approach that will be useful for the analysis of classical bifurcations. We investigate the existence, uniqueness and blow up of the solutions and show that noise can prevent explosion. The stochastic stability of the singular point is characterized and the absorption time at the singular point is shown to be almost surely finite if and only if $\alpha < 1$. In that case, we show that quasi-stationary distributions exist when $\alpha < \frac{3}{4}$. In the case $\alpha \geq 1$, we study the existence of other stationary distributions. Beyond the mathematical developments, this paper is also resolutely intended for engineers, physicists and other scientists interested in the qualitative behavior of noisy systems described through stochastic differential equations. Therefore, readers interested in applications may skip this section and jump directly to section 3 where they will find readily applicable results. In that section we build on the results of the general analysis to analyze the stochastic pitchfork, Hopf and saddle-node bifurcations with power diffusion function. Interestingly, in all the cases, any system locally reducible around an equilibrium point to one of these normal forms displays the same qualitative behavior, provided that the drift satisfies an additional condition at infinity. This property is analogous to the well-known universal unfolding property for deterministic dynamical systems. In all three cases, we put in good use the characterization of the dynamics by developing applications to the Wilson and Cowan system [19], a simple neural population model widely used in neurosciences.


noise vanishes at 0
Noise-induced stabilization of the saddle point
Fluctuations around the stable equilibrium

Figure 1: Stability of a singular saddle point in double-well potential perturbed by power diffusion coefficient. A particle located at the saddle point remains steady. If a small perturbation is applied, two scenarii are possible according to the local behavior of $\gamma(x)$ (see theorem 1): either noise brings back the particle to the saddle point (left) or the particle falls into the well and fluctuates without ever reaching the saddle point (right).

2. General Theory

In this section we analyze the behavior of the solutions of a general equation (1) under a few assumptions on the function g and γ:

(H1). $g(x) \sim \mu x^{1+\kappa}$ at $x = 0$, for some $\mu \in \mathbb{R}$ and $\kappa > 0$

(H2). $g(x) \leq -\nu x^{1+\beta}$ when $|x| \geq A$ for some $A > 0$, for $\nu \in \mathbb{R}$ (generally assumed non-negative) and $\beta > 0$.

(H3). $\gamma(x) \sim \sigma |x|^{\alpha}$ at $x = 0$, for some $\sigma \geq 0$ and $\alpha \geq 0$

(H4). $\gamma(x) \sim d\, x^{\delta}$ at $x = \pm\infty$ with $d > 0$ and $\delta \geq 0$.

(H5). $\gamma(x)$ only vanishes at $x = 0$

(H6). Either (i) $\nu \geq 0$ or (ii) $\nu < 0$ and $\delta > 1 + \beta/2$.

Assumptions (H1) and (H3) describe the behavior of the solutions close from zero and will govern the stability of the singular point 0. The assumptions (H2), (H4) and (H5) are global properties that will be chiefly used for explosion matters, and existence of stationary distributions. Such assumptions are generally not necessary in the deterministic case: indeed, in the local unfolding of a bifurcation, the presence of a non-zero fixed point implies the local existence of a manifold of fixed points under some non-degeneracy conditions. However, in the stochastic case, as soon as the singular point (where noise vanishes) is unstable, the system will visit very regions very remote from the singular point with positive probability, and these latter conditions will come into play.

The stochastic process distributed as a Dirac at zero ($X_t = 0$ a.s.) is solution of the equation whatever the parameters. Studying the stationary solutions and their stability will make essential use of the differential operator $\mathcal{L}$ associated with Itô's representation of the diffusion acting on twice differentiable functions $V \in C^2(\mathbb{R}, \mathbb{R})$:

$$\mathcal{L}V(x) = (\lambda x + g(x))\, V'(x) + \frac{1}{2}\gamma(x)^2 V''(x).$$

Beyond the well-posedness of the equation studied below in section 2.1, the main results of the present section can be summarized in the following:

Theorem 1. *Under the assumptions (H1)- (H5) we have:*

- *for* $\frac{1}{2} \leq \alpha < 1$*, the singular point is stable in probability and any solution is absorbed in finite time almost surely at the singular point. If moreover* $\alpha < 3/4$*, there exists an exponentially attractive quasi-stationary distribution for the trajectories that did not reach the singular point.*
- *for* $\alpha = 1$*, the singular point is stable for* $\lambda < \sigma^2/2$ *and unstable otherwise, and two stationary solutions exist for* $\lambda > \sigma^2/2$*.*
- *for* $\alpha > 1$*, the singular point is stable for* $\lambda < 0$ *and unstable otherwise, and two stationary solutions exist for* $\lambda > 0$*.*

2.1. Existence and uniqueness of solutions

The first question arising with equation (1) is its well-posedness. Indeed, this equation does not satisfy the standard conditions for existence and uniqueness of solutions: the drift function is neither globally Lipschitz-continuous, nor holds a linear growth condition. Moreover, the diffusion coefficient is also singular: for $\alpha < 1$, the diffusion coefficient is not Lipschitz continuous at 0, and for $\alpha > 1$ it does not satisfies the linear growth condition generally imposed for the existence and uniqueness. Based on existing results [20], we prove the following property:

Proposition 1. *For any* $\alpha \in [\frac{1}{2}, \infty)$*, there exists a unique strong solution to the equation* (1)*. Under the assumptions (H2), (H4) and (H6), the solution never blows up in finite time.*

Proof. In the case where $\alpha \geq 1$ it is clear that both the drift and diffusion functions are locally Lipschitz continuous. This property implies that the strong uniqueness property of the solutions holds (see e.g. [20, Chap. IX.2]). Moreover, the drift and the diffusion functions are locally bounded, ensuring the existence of solutions up to an explosion time. These two properties ensure strong existence and uniqueness of solutions up to an explosion time.

The case $\alpha < 1$ is slightly more complex. In that case, the linear growth condition is clearly satisfied, but the diffusion coefficient is not Lipschitz at 0. However, the method presented in [20, theorem IX.3.5] based on the properties of the local time at zero of the diffusions ensure that we have the existence and uniqueness of the solutions of (1) for $\alpha \geq \frac{1}{2}$ and non-uniqueness of solutions for $\alpha < \frac{1}{2}$.

Let us now address the non-explosion property. To this purpose, we use Feller's test for explosion (see [21]). The function $\frac{\lambda x + g(x) + 1}{\gamma(x)^2}$ is locally integrable on both the intervals $(-\infty, 0)$ and $(0, \infty)$. We consider the interval $(0, \infty)$ for instance (the same principle applies for the interval $(-\infty, 0)$), and for some $c > 0$, we define the scale function:

$$p(x) := \int_c^x \exp\left(-2\int_c^y \frac{\lambda\xi + g(\xi)}{\gamma^2(\xi)}\, d\xi\right) dy.$$

The process almost surely never diverges before possibly reaching zero as soon as soon as $p(x)$ is diverges at ∞. At infinity, we have:

$$-2\int_c^y \frac{\lambda\xi + g(\xi)}{\gamma^2(\xi)} \geq (\frac{2\nu}{d^2} - \varepsilon)x^{2+\beta-2\delta}$$

for some $\varepsilon > 0$, arbitrarily small when x is sufficiently large. This quantity is lower bounded at infinity in the case where $\nu > 0$ or in the case $\delta > 1 + \beta/2$, implying in both case that $p(x) \to \infty$ when $x \to \infty$. We hence conclude that the process does not blow up in finite time in these cases, which ends the proof. ☐

Remark. *This property illustrates a first stabilization effect of power diffusions, namely the blow-up prevention. In the case where the solutions to the noiseless equation do not blow up in finite time, power diffusion coefficients cannot create a stochastic blow up. In the case where the solutions to the noiseless system blow up in finite, a power diffusion function prevents blow up provided that the diffusion coefficient diverges fast enough at infinity compared to the diffusion function.*

2.2. Stability of δ_0

We now address the stability of the equilibrium δ_0, and prove the following:

Theorem 2. *Let us denote by δ_0 the stationary solution $X_t = 0$ a.s. for all t. We have:*

- *For $\alpha < 1$, δ_0 is asymptotically stable in probability whatever $\lambda \in \mathbb{R}$ and $\sigma \neq 0$.*
- *For $\alpha = 1$, δ_0 is:*
  - (i). *asymptotically stable* in probability *for $\lambda < \frac{\sigma^2}{2}$. Moreover, if $g(x) = -\mu x^{1+\kappa}$ for $\mu, \kappa > 0$ and $\gamma(x) = \sigma x$, the stationary solution δ_0 is* almost surely exponentially stable *for $\lambda < \frac{\sigma^2}{2}$.*
  - (ii). *unstable in probability for $\lambda > \frac{\sigma^2}{2}$*
- *For $\alpha > 1$, δ_0 is :*
  - *stable in probability for $\lambda < 0$*
  - *unstable in probability for $\lambda > 0$*

Proof. **Case** $\alpha < 1$: Let us consider $V(x) = x^\alpha$ as a Lyapunov functional for the dynamics. This function is clearly $C_2^0(\mathbb{R})$, i.e. it is twice continuously differentiable except at $x = 0$. Moreover, we have:

$$\mathcal{L}V = \alpha x^{\alpha-1}(\lambda x + g(x)) - \frac{\alpha(1-\alpha)}{2}\,\gamma(x)^2 x^{\alpha-2}$$

Since we assume that $\alpha < 1$ and $x = o(g(x))$ at 0, the leading term close to 0 is of order $-\alpha(1-\alpha)/2\ \sigma^2 x^{3\alpha-2}$ which is strictly negative for sufficiently small x. More precisely, there exists $r > 0$ such that $V \in C_2^0(U_r)$ where U_r is the open ball of radius r and such that $\mathcal{L}V < 0$ for all $x \in U_r$. Moreover, for any ε such that $0 < \varepsilon < r$ we have $V(r) > 0$ and $\mathcal{L}V < -c_\varepsilon < 0$ for all $r > x > \varepsilon$. Theorem V.4.1 of [17] therefore applies and concludes the proof of the stability in probability of the solution δ_0 whatever the parameters $\lambda \in \mathbb{R}$, $\sigma > 0$ and $\alpha \in [1/2, 1)$.

Case $\alpha = 1$: We start considering the case $\lambda < \sigma^2/2$ and define $\eta = \frac{1}{2} - \frac{\lambda}{\sigma^2} > 0$. The Lyapunov function $V : x \mapsto x^\eta$ satisfies:

$$\mathcal{L}V\,(x) = \eta x^\eta(\lambda + \frac{1}{2}(\eta - 1)\gamma^2(x)) + \eta x^{\eta-1}\,g(x).$$

Under assumption (H1) we have $x\,g(x) \sim \mu x^{2+\kappa}$, hence is equivalent, close from zero, to

$$\eta x^\eta \left(\lambda + \frac{(\eta-1)}{2}\frac{\gamma^2(x)}{x^2}\right) = \frac{1}{2}(\lambda - \frac{\sigma^2}{2})\eta x^\eta.$$

Since $(\lambda - \sigma^2/2) < 0$, the same argument as in the case $\alpha < 1$ ensures asymptotic stability of δ_0 in probability for $\lambda < \sigma^2/2$. If we now further assume $g(x) = -\mu x^{1+\kappa}$ and $\gamma(x) = \sigma\,x$, considering $V(x) = x^2$, we have:

- $\mathcal{L}V\,(x) = (2\lambda + \sigma^2)x^2 - 2\,\mu\,x^{2+\kappa} \leq (2\lambda + \sigma^2)V(x)$,
- $|V'(x)\sigma\,x|^2 = |2\sigma x^2|^2 = 4\sigma^2 V(x)$

Using the stochastic stability theorem [18, Theorem 3.3], we deduce that:

$$\limsup_{t\to\infty} \frac{1}{t}\log|X(t)| \leq -\frac{4\sigma^2 - 2(2\lambda + \sigma^2)}{4} = \lambda - \frac{\sigma^2}{2} \qquad \text{almost surely.}$$

By definition of the almost sure exponential stability, we conclude that the solution 0 is almost surely exponentially stable for any $\lambda < \frac{\sigma^2}{2}$.

Let us now deal with the case $\lambda > \sigma^2/2$. We define $V(x) = -\log(x)$. This function tends to infinity as $x \to 0$ and belongs to $C_2^0(\mathbb{R})$. Moreover,

$$\mathcal{L}V = (-\lambda + \frac{\gamma(x)^2}{2x}) + \frac{g(x)}{x}.$$

which is equivalent at zero to $-\lambda + \sigma^2/2 < 0$. Hence there exists $r > 0$ such that $\mathcal{L}V \leq 0$ for any $x < r$, and moreover we have for any $\varepsilon > 0$ and $r > x > \varepsilon$, $V(x) \geq 0$ and $\mathcal{L}V(x) < -c_\varepsilon < 0$. Theorem V.4.2 of [17] applies and ensures that the solution δ_0 is not stable in probability.

Case $\alpha > 1$**:** For $V(x) = x^2$ around zero, we have:

$$\mathcal{L}V = 2\lambda x^2 + 2xg(x) + \gamma(x)^2$$

and under the assumption that $g(x) \sim \mu x^{1+\kappa}$ at zero and $\alpha > 1$, the term $\mathcal{L}V$ is equivalent to $2\lambda x^2$ and hence locally has the sign of λ. For $\lambda < 0$, we can directly apply theorem V.4.1 of [17]. For $\lambda > 0$, we use $V(x) = -\log(x)$ again and conclude that 0 is not stable in probability. □

2.3. Extinction time

We know that $X_t = 0$ a.s. is a stationary solution of the SDE. It is an absorbing state of the diffusion, i.e. as soon as the solution reaches the singular point, the solution is almost surely identically stuck at the singular point for the whole subsequent evolution. We are now interested in the extinction probability, namely the probability of reaching zero in finite time. We distinguish between the cases $\alpha < 1$ and $\alpha \geq 1$.

2.3.1. Case $\alpha < 1$

Proposition 2. *Under assumptions (H1)-(H6), the first hitting time of zero of the solution of*

$$dx_t = (\lambda x + g(x))\,dt + \gamma(x)dW_t$$

is almost surely finite for any $\alpha < 1$ and $\sigma > 0$. This is also the case for $\nu = 0$ and $\lambda < 0$.

Proof. In order to establish this property, we use a sufficient condition derived from Feller's test for expositions and evidence the almost sure finite exit time of the interval $]0,\infty[$. The finiteness of the exit time of $]0,\infty[$ will readily imply the finiteness of the hitting time of 0, since the process is almost surely bounded for all finite time as proved in proposition 1 under the assumptions of the proposition.

Let us define the interval $I = (0,\infty)$, and let $c \in I$ arbitrarily chosen. Under assumption (H5), we have that for all $x \in I$, the diffusion coefficient is always strictly positive, hence the function $(1+|\lambda x+g(x)|)/\gamma(x)^2$ is locally integrable on I. The finiteness of the exit time of the process from the interval I can be established through the analysis of Feller's scale functions. For convenience, we introduce the function:

$$G(x) = \int_c^x \frac{(\lambda\xi + g(\xi))}{\gamma^2(\xi)} d\xi.$$

Feller's scale functions written as a function of G read:

$$\begin{cases} p(x) & \overset{def}{=} \int_c^x \exp\{-2G(y)\}\, dy \\ m(dx) & \overset{def}{=} \frac{2\,\exp(2\,G(x))}{\gamma^2(x)}\, dx \\ v(x) & \overset{def}{=} \int_c^x (p(x) - p(y))\, m(dy) \end{cases}$$

We prove that (a) $\lim_{x\to\infty} p(x) = \infty$ and (b) $\lim_{x\to 0^+} v(x) < \infty$. The property (a) has been addressed in the proof of proposition 1. We hence only need to prove the property (b) which is slightly more subtle. The function $v(x)$ is indeed defined as an integral, which is well behaved except possibly at $x = 0$. Close to this point, we have:

$$\begin{aligned} G(y) - G(z) &= \int_z^y \frac{\lambda\xi + g(\xi)}{\gamma^2(\xi)} d\xi \\ &\sim_{0^+} \frac{\lambda}{2\sigma^2(1-\alpha)}(y^{2(1-\alpha)} - z^{2(1-\alpha)}) + \frac{\mu}{\sigma^2\phi}(y^\phi - z^\phi) \end{aligned}$$

with $\phi = 2 + \kappa - 2\alpha$ and this function is smooth and bounded close to zero. Let us denote by M the supremum of $|G(y) - G(z)|$ on $[0, c]$, and by $F(x, y)$ the function defined for $0 < y \leq x \leq c$:

$$F(x,y) = \int_y^x \exp(2(G(y) - G(z)))\, dz \leq e^{2M}(x-y).$$

It is then clear that:

$$v(x) = 2\int_c^x \frac{F(x,y)}{\gamma^2(y)}\, dy \leq 2\int_c^x \frac{e^{2M}(x-y)}{\gamma^2(y)}\, dy$$

which is integrable at zero since the integrand is of order $x^{1-2\alpha}$ and hence integrable at zero under the condition that $\alpha < 1$. By application of Feller's test for explosion [21, Prop. 5.32 (iii)] we conclude that if S denotes the exit time of the process from the interval I, we have $\mathbb{P}(S < \infty) = 1$.

Moreover, we have seen in proposition 1 that under the assumptions of the proposition, the process is almost surely bounded. We hence conclude that necessarily the process reaches 0 in finite time and is absorbed (this is also a direct consequence of Feller's test for explosion [21, Prop. 5.22] under the condition that $p(x) \to \infty$ when $x \to \infty$ if $p(x) > -\infty$ when $x \to 0^+$, which is a simple consequence of the above estimates).

Let us now deal with the case $\nu = 0$ and $\lambda < 0$, we have $G(x) = \lambda x^{2(1-\alpha)} + C$ for some constant C. The same reasoning as above ensures that $p(x)$ tends to infinity as $x \to \infty$. Demonstrating that $\nu(x)$ is bounded at zero did not use the fact that $\nu \neq 0$ and hence applies in the present case. We conclude that $P(S < \infty) = 1$, hence the process almost surely reaches 0 in finite time.

□

We hence proved that for $\alpha < 1$, the solutions almost surely hit zero in finite time, where they stay trapped for the subsequent dynamics, as soon as the diffusion coefficient σ is nonzero.

When $\sigma = 0$, this is never the case: in the case $\lambda < 0$, the solutions of the related ODE converge exponentially fast towards zero but never actually reach it, and in the case $\lambda > 0$, the solutions escape from zero exponentially. It is hence of interest to analyze the behavior of the system in the small noise limit. We focus on the case $\lambda > 0$ (in which case 0 is unstable in the noiseless system and a.s. reached in finite time in the stochastic case) and use Freidlin and Wentzell large deviations theory [22] to analyze in detail the behavior of the system in the small noise limit. In order to obtain precise quantitative estimates, we will assume that $g(x) = -\mu x^{1+\kappa}$ with $\kappa > 0$ and $\mu > 0$ and $\gamma(x) = \sigma |x|^{\alpha}$, corresponding to a double-well potential. Note that the method developed here does not depend on the choice of $g(x)$.

We analyze the properties of the extinction time of the system with initial condition chosen as the fixed point of the noiseless system:

$$\tau_0 := \inf \left\{ t > 0;\ X_t = 0 \quad | \quad X_0 = \left(\frac{\lambda}{\mu}\right)^{1/\kappa} \right\}$$

In the case $\alpha < 1$, one can expect from the large deviation theory that $\mathbb{E}[\tau_0]$ becomes exponentially large as $\sigma \to 0$, i.e behaves in the leading order as e^{V/σ^2} where V is a positive constant. More precisely, we have:

Proposition 3. *For all* $\frac{1}{2} < \alpha < 1$ *and* $\lambda > 0$*:*

$$\lim_{\sigma \to 0} \sigma^2 \log \mathbb{E}[\tau_0] = \lambda \left(\frac{\lambda}{\mu}\right)^{2(1-\alpha)/\kappa} \left[\frac{1}{2(1-\alpha)} - \frac{1}{2(1-\alpha)+\kappa} \right] > 0$$

Proof. By application [22, Theorem 4.1], the problem is reduced to compute at the boundary $x = 0$ the quasipotential $U(x)$, solution of the Hamilton-Jacobi equation:

$$(\lambda x - \mu x^{1+\kappa}) U'(x) + \frac{x^{2\alpha}}{2} (U'(x))^2 = 0 \tag{2}$$

with boundary condition $U((\frac{\lambda}{\mu})^{\frac{1}{\kappa}}) = 0$. That is, one has to compute:

$$U(x) = \int_x^{\left(\frac{\lambda}{\mu}\right)^{1/\kappa}} \frac{\lambda u - \mu u^{1+\kappa}}{u^{2\alpha}} du \tag{3}$$

Simple algebra yields the expression:

$$U(0) = \lambda \left(\frac{\lambda}{\mu}\right)^{2(1-\alpha)/\kappa} \left[\frac{1}{2(1-\alpha)} - \frac{1}{2(1-\alpha)+\kappa}\right] \tag{4}$$

□

As a consequence, for $\alpha < 1$ fixed, the extinction time becomes exponentially large, which explains the transition between the regime $\sigma > 0$ (where the system a.s. hits zero) and $\sigma = 0$. A critical boundary of this domain is the point $\alpha = 1$: in that case, the system a.s. never hits zero in finite time as shown in section 2.3.2.

In particular, proposition 3 ensures that the convergence as $\alpha \to 1^-$ with $\lambda > 0$ fixed displays a super-exponential time behavior:

$$\lim_{\alpha\to 1^-} \lim_{\sigma\to 0} \quad \sigma^2 \log \mathbb{E}\left[\tau_0\right] = \infty$$

Going deeper into the analysis of the above convergence allows to understand more precisely what happens in the critical parameter region where both $\alpha \to 1^-$ and $\lambda \to 0^+$, in the limit of small noise $\sigma \to 0$. Such a "blow up" reveals the following picture around the point $(\alpha, \lambda, \sigma) = (1, 0, 0)$ in parameter space:

Corollary 1. *Assume that $\lambda \to 0^+$ and $\alpha \to 1^-$ such that*

$$\frac{\lambda}{(1-\alpha)} \to c \in [0, \infty] \tag{5}$$

Then, one has the following three regimes:

- *if $c = 0$, the extinction time is sub-exponential :*

$$\lim_{\alpha\to 1, \lambda\to 0} \lim_{\sigma\to 0} \quad \sigma^2 \log \mathbb{E}\left[\tau_0\right] = 0$$

- *if $0 < c < \infty$, the extinction time is exponential :*

$$\lim_{\alpha\to 1, \lambda\to 0} \lim_{\sigma\to 0} \quad \sigma^2 \log \mathbb{E}\left[\tau_0\right] = \frac{c}{2\mu^{2(1-\alpha)/\kappa}}$$

- *if $c = \infty$, the extinction time is super-exponential :*

$$\lim_{\alpha\to 1, \lambda\to 0} \lim_{\sigma\to 0} \quad \sigma^2 \log \mathbb{E}\left[\tau_0\right] = \infty$$

where $\lim\limits_{\alpha\to 1, \lambda\to 0}$ is constrained by Eq. (5).

Proof. This proposition is a consequence of the analysis of the different asymptotic regimes of the critical exponent c^* obtained in proposition 3:

$$c^* = \lambda \left(\frac{\lambda}{\mu}\right)^{2(1-\alpha)/\kappa} \left[\frac{1}{2(1-\alpha)} - \frac{1}{2(1-\alpha)+\kappa}\right].$$

The leading term in the considered limit is proportional to $\lambda^{1+2(1-\alpha)/\kappa}/(1-\alpha)$. A series expansion provides the following expression:

$$c^* \sim \frac{1}{2\mu^{2(1-\alpha)/\kappa}}(\frac{\lambda}{\alpha-1}+\frac{2}{\kappa}\lambda\log(\lambda))+O(\lambda\log(\lambda)^2(1-\alpha)).$$

The different regimes identified in the corollary immediately follow from the analysis of the limits of this exponent as $\lambda \to 0^+$ and $\alpha \to 1^-$ subject to the condition (5). $\square$

Remark. *Note that the regimes do not depend on μ or κ qualitatively and the property is purely local: as λ goes to zero, the integration range becomes increasingly small. The property is hence valid for equation* (1) *with function $g(x)$ equivalent at zero to $-\mu x^{1+\kappa}$. However, the asymptotic exponential divergence rate quantitatively depends on the local behavior of the flow close to zero, as observed in the case $0 < c < \infty$.*

From a dynamical systems viewpoint, this type of solutions corresponds to a relatively sharp convergence towards the equilibrium. Indeed, the dynamics stops at the almost sure hitting time of the singular point and the whole subsequent dynamics is deterministic and constant. This will not be the case for more regular diffusion coefficients ($\alpha \geq 1$) as we now show.

2.3.2. *For $\alpha \geq 1$ the solution a.s. never reaches* 0

We now deal with the case $\alpha \geq 1$ and show that in contrast to the case $\alpha < 1$, even when δ_0 is stable in probability or almost surely exponentially stable, the solutions of the SDE almost surely never reaches zero.

Theorem 3. *Let $\alpha \geq 1$. Then for any $x_0 \in \mathbb{R}$ we have*

$$\mathbb{P}[x_t \neq 0 \text{ on } t \geq 0] = 1$$

That is, almost any sample path starting from a non-zero initial condition will never reach zero.

Remark. *A similar result (for $\alpha = 1$) in proved for instance in [18, lemma 3.2].*

Proof. Denote by τ the first hitting time of zero of the process with initial condition $x_0 \neq 0$ at $t = 0$. Assuming that the conclusion is false, we can find a pair of constants $T > 0$ and $\theta > 1$ sufficiently large such that $P(B) > 0$ where B is defined as:

$$B = \{\tau \leq T \text{ and } |x(t)| \leq \theta - 1 \text{ for all } t \leq \tau\}$$

Moreover, under the assumption $\alpha \geq 1$, both the drift and diffusion functions are locally Lipschitz-continuous, hence there exists a positive constant K_θ such that:

$$|\lambda x + g(x)| \vee \gamma(x) \leq K_\theta |x| \quad \text{for all } |x| \leq \theta,\ t \leq T.$$

Let now $V(x) := |x|^{-1}$. We have for $0 < |x| \leq \theta$ and $t \leq T$

$$\mathcal{L}V \leq K_\theta(1+K_\theta)V(x)$$

Let us now define $\tau_\varepsilon = \inf\{t \geq 0;\ |x_t| \notin (\varepsilon, \theta)\}$. This is a stopping time, so by Itô's formula and the optimal sampling theorem with the bounded stopping time $\tau_\varepsilon \wedge T$, we have:

$$\begin{aligned}\mathbb{E}[e^{-K_\theta(1+K_\theta)(\tau_\varepsilon\wedge T)}V(x_{\tau_{\varepsilon\wedge T}})] &= V(x_0)+\\ &+\mathbb{E}\Big[\int_0^{\tau_\varepsilon\wedge T} e^{-K_\theta(1+K_\theta)\,s}[-K_\theta(1+K_\theta)V(x_s)+(\mathcal{L}V)(x_s)]\,ds\Big]\\ &\leq |x_0|^{-1}\end{aligned}$$

We know that for any $\omega \in B$, the time τ_ε is smaller than T and moreover that $x_{\tau_\varepsilon} = \varepsilon$ by definition of B, hence we have:

$$|x_0|^{-1} \geq \mathbb{E}[e^{-K_\theta(1+K_\theta)(\tau_\varepsilon \wedge T)} V(x_{\tau_{\varepsilon \wedge T}})] \geq \mathbb{E}\Big[e^{-K_\theta(1+K_\theta)T} \varepsilon^{-1} \mathbb{1}_B\Big]$$

i.e.

$$\mathbb{P}(B) \leq \varepsilon |x_0|^{-1} e^{K_\theta(1+K_\theta)T}$$

This inequality is true whatever ε (we recall that the definition of B did not involve ε), hence valid in particular when ε is arbitrarily small, and necessarily $\mathbb{P}(B) = 0$ which contradicts the definition of B. □

2.4. Stationary and quasi-stationary solutions

We are now interested in identifying possible stationary solutions. We already mentioned that the null stochastic process (distributed as a Dirac delta measure at 0) is always a stationary solution of the system. Moreover, for $\alpha < 1$, we have shown that the process is almost surely absorbed at zero in finite time. This property prevents from the existence of any stationary solution, and as far as the permanent dynamics is concerned, it is perfectly characterized by the absorption property. In order to further describe the behavior of the solutions, one may wonder how the solutions that did not yet reach zero are distributed. Stationary solutions of the process conditioned on never hitting zero are called quasi-stationary distributions. In the case $\gamma(x) = \sigma|x|^\alpha$ with $\alpha < 3/4$, we can show that there exists a quasi-stationary distribution that attracts exponentially fast the conditioned process. In detail, we have:

Proposition 4. *Let us assume that $v > 0$, g is differentiable, g' is bounded at zero and smaller than $Cx^{2\beta}$ for some $C \in \mathbb{R}$ at infinity, that assumptions (H1)-(H6) hold and $\alpha \in [1/2, 3/4)$. The process conditioned on never reaching zero and with a bounded initial distribution converges exponentially fast towards a probability measure $p_\pm$ for any positive (resp. negative) initial condition. These are called quasi-stationary solutions, or Yaglom limit.*

Proof. We apply a result due to Cattiaux and collaborators [15] in the domain of diffusion models in population dynamics. In their article, they prove consider a diffusion of type:

$$dX_t = dB_t - q(X_t)\, dt$$

under a few conditions on q. Our problem can be brought to their framework changing variables and considering: $X_t = x_t^{1-\alpha}/(\sigma(1-\alpha))$. Itô's formula allows putting our equations in that form, with

$$q(x) = -\frac{1}{\sigma} Z^{-\alpha} f(Z) + \frac{\alpha}{2(1-\alpha)} \frac{1}{x}$$

and where $f(x) = \lambda x + g(x)$ and $Z = (\sigma(1-\alpha)x)^{1/(1-\alpha)}$. Under the conditions that (i) the first hitting time of zero is almost surely finite and (ii) that the conditions: $\Delta = -\inf_{y \in (0,\infty)} q^2(y) - q'(y) < \infty$ and $\lim_{y \to \infty} q^2(y) - q'(y) = \infty$, they show that the spectrum of the Kolmogorov backward equation (for the related operator $\mathcal{L}$) has a purely discrete spectrum with no zero eigenvalue. Moreover, they show that a rescaled version of the related eigenfunction is integrable and exponentially attract all initial conditions under the conditions:

$$\int_0^1 \frac{1}{q(y)^2 - q'(y) + \Delta + 2} e^{-G(y)}\, dy < \infty \tag{6}$$

with $G(y) = \int_1^x 2q(y)\,dy$,

$$\int_1^\infty e^{-G(y)}dy < \infty \ , \ \int_0^1 xe^{-G(x)/2}dx < \infty \text{ and } \int_1^\infty e^{G(y)} \int_y^\infty e^{G(z)}\,dzdy < \infty \tag{7}$$

In order to apply their result, we demonstrate that all these conditions are satisfied in our framework. First of all, under the condition $\alpha < 1$ and $\nu > 0$, we showed that the process x_t solution of (1) almost surely reach zero in finite time (proposition 2). This is hence also the case of X_t since $(1 - \alpha) > 0$ in that case. Moreover, the derivative of the function $q(x)$ reads:

$$q'(x) = \alpha Z^{-1} f(Z) - f'(Z) - \frac{\alpha}{2(1-\alpha)}\frac{1}{x^2}.$$

Around zero, because of our assumptions on g, we have:

$$q(x) \sim \frac{\alpha}{2(1-\alpha)}\frac{1}{x} \text{ and } q'(x) \sim -\frac{\alpha}{2(1-\alpha)}\frac{1}{x^2}$$

and hence

$$q^2(x) - q'(x) \sim_{x\to 0^+} \frac{\alpha(2-\alpha)}{4(1-\alpha)^2}\frac{1}{x^2}$$

which tends to infinity as $x \to 0^+$. At $x = \infty$, we have:

$$q^2(x) - q'(x) \geq \frac{\mu^2}{\sigma^2}(1-\varepsilon)Z^{2(1-\alpha)+2\beta}$$

for any $\varepsilon > 0$ and hence tends to infinity when $x \to \infty$. In particular, $\Delta = -\inf_{y\in(0,\infty)} q^2(y) - q'(y) < \infty$

Let $G(x) = \int_1^x 2\,q(y)\,dy$. We now show that the condition (6) is satisfied. At infinity, $q(x)$ is greater or equal to $C' x^{1+\beta/(1-\alpha)}$ with $C' > 0$. It is then easy to show that $\lim_{x\to\infty} \frac{G(x)}{x} = \infty$. Moreover, we have

$$A := \lim_{x\to 0^+} \left(G(x) - \frac{\alpha}{1-\alpha}\log(x)\right) \in (-\infty, \infty).$$

Therefore, near zero, the integrand involved in the expression of $Q(x)$ is proportional to $y^{2-\frac{\alpha}{1-\alpha}}$. Under the condition $\alpha < \frac{3}{4}$, condition (6) holds. Showing that condition (7) is valid is straightforward at this point: G grows at least linearly at ∞ ensuring that $\int_1^\infty e^{-G(y)}dy < \infty$. Moreover, around zero, the function $xe^{-G(x)/2}$ is of order $x^{1-\frac{1}{2}\frac{\alpha}{1-\alpha}}$ which is integrable at zero for $\alpha < \frac{4}{5}$, hence in particular for $\alpha < \frac{3}{4}$. The main results of [15] (namely their theorem 5.2, proposition 5.5, corollaries 6.1 and 6.2) apply, and ensure the existence and uniqueness of a quasi-stationary distribution attracting exponentially fast the law of the diffusion conditioned on not hitting zero with initial laws having bounded support. □

Let us now deal with the general case with diffusion coefficient $\gamma(x)$ and $\alpha \geq 1$. In that case, the process is almost surely never absorbed at zero. Possible stationary solutions with smooth probability density function p with respect to Lebesgue's measure satisfy the forward Kolmogorov (or Fokker-Planck) equation:

$$\frac{\mathrm{d}}{\mathrm{d}x}\left((\lambda\, x + g(x))p(x) - \frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}x}\left(\gamma(x)^2 p(x)\right)\right) = 0. \tag{8}$$

i.e.
$$(\lambda\, x + g(x) - \gamma'(x)\gamma(x))p(x) - \frac{1}{2}\gamma(x)^2 p'(x) = K$$
for some $K \in \mathbb{R}$. If we can find solutions to these equations subject to the conditions $p(x) \geq 0$ for all $x \in \mathbb{R}$ and $\int_{\mathbb{R}} p(x)\, dx = 1$ (*acceptable solutions*), then these are probability density functions of stationary solutions of the diffusion equation. For simplicity, we consider that assumption (H5) is valid, i.e. $\gamma(x)$ only vanishes at $x = 0$. In that case, since δ_0 is a solution to the SDE (1), stationary solutions will either have support on $\mathbb{R}^+$ or $\mathbb{R}^-$. In the case where the dynamics has additional singular points, the Kolmogorov equation will define solutions in each interval delimited by the singular points (see section 3.3). The Kolmogorov equation is a linear ordinary differential equation with constant source term. The homogeneous solutions (with $K = 0$) is given by:
$$p_0(x) = C\gamma(x)^{-2} \exp\left(\int_{.}^{x} (\frac{2\lambda y}{\gamma(y)^2} + 2\frac{g(y)}{\gamma(y)^2})\, dy\right)$$
defined for $x \neq 0$. Particular solutions are given by:
$$p_K(x) = \left(-2K \int_1^x \exp\left(-2\int_1^y \frac{2\lambda z}{\gamma(z)^2} + 2\frac{g(z)}{\gamma(z)^2}\, dz\right) dy + \gamma(1)^2 p(1)\right) \times \gamma(x)^{-2} \exp\left(\int_1^x (\frac{2\lambda y}{\gamma(y)^2} + 2\frac{g(y)}{\gamma(y)^2})\, dy\right) \quad (9)$$

We have the following:

Proposition 5. *In addition to the stationary distribution δ_0, we have:*

- *For $\alpha = 1$ and:*
  - $\lambda < \sigma^2/2$ *there exists no integrable solution of the Kolmogorov equation*
  - $\lambda \geq \sigma^2/2$, *under assumption (H6), there exist two integrable solutions of the Kolmogorov equation given by $Z_+ p_0(x)\mathbb{1}_{x>0}$ and $Z_-\, p_0(x)\mathbb{1}_{x<0}$ where Z^+ and Z^- generically denote normalization constants.*
  - $\lambda \geq \sigma^2/2$, $\nu < 0$ *and* $\delta < 1 + \frac{\beta}{2}$ *(case where assumption (H6) is not satisfied) there does not exist any stationary solution (we saw that the solutions a.s. blow up in finite time)*
- *For $\alpha > 1$ and:*
  - $\lambda < 0$ *the Kolmogorov equation has no integrable solution*
  - $\lambda > 0$ *and under assumption (H6), the Kolmogorov equation has two integrable solution given by $Z_+\, p_0(x)\mathbb{1}_{x>0}$ and $Z_-\, p_0(x)\mathbb{1}_{x<0}$.*
  - $\lambda > 0$, $\nu < 0$ *and* $\delta \leq 1 + \frac{\beta}{2}$ *(i.e. when (H6) is not satisfied), the Kolmogorov equation has no integrable solution.*

Remark. *In the case $\alpha < 1$, we already mentioned that no stationary solution can exist. We can show using the expressions of the solutions to the Kolmogorov equation that no solution is integrable.*

Proof. For $\alpha > 1$, let us analyze the integrability of $p_0(x)$. Around 0, the function $x^{-2\alpha}$ is not integrable. The leading term of the exponential around zero is given by $\frac{\lambda}{\sigma^2(1-\alpha)} x^{2(1-\alpha)}$, hence diverges at zero. If $\lambda < 0$, this term tends to $+\infty$ and is not integrable, and if $\lambda > 0$, the exponential term tends to zero and the function is hence integrable at 0 and $p(0) = 0$. For $x \to \infty$, we distinguish two cases:

- for $\delta < 1 + \beta/2$ and $\nu > 0$, the leading term of the exponential in the expression of $p_0(x)$ is given by $-\nu z^{1+\beta-2\delta}/(1+\beta-2\delta)$ which tends to $-\infty$ for $\nu > 0$ fast enough ensuring integrability of $p_0(x)$ at ∞. For $\nu < 0$, this term tends to ∞ and $p_0(x)$ is not integrable at infinity.
- when $\nu < 0$, the leading term of the exponential at infinity is of order $-\nu z^{1+\beta-2\delta}/(1+\beta-2\delta)$ which tends to $+\infty$ and hence the solution to the Kolmogorov equation is not integrable at infinity.
- For $\delta \geq 1 + \beta/2$, the terms inside the exponential tend to zero, and the integrability relies on the integrability property of $x^{-2\delta}$, always integrable at infinity.

We hence proved that acceptable solutions related to $K = 0$ exist for $\lambda > 0$ and no solution existed when $\lambda < 0$. Moreover, there is no possible choice of $K \neq 0$ that can overcome the exponential divergence at zero. Indeed, the only possible choice of K would be the only constant preventing divergence at zero, i.e.:

$$p(1) = 2\frac{K}{\sigma^2} \int_1^0 \exp\left(-2\int_1^y \frac{2\lambda z}{\gamma(z)^2} + 2\frac{g(z)}{\gamma(z)^2}\,dz\right) dy.$$

However, the integral term of the righthand side diverges, hence the above relationship cannot be satisfied and there is no acceptable solution for $\lambda < 0$. Eventually let us remark that for $\lambda > 0$, any choice of K prevents integrability at infinity.

For $\alpha = 1$, we again analyze the integrability properties of $p_0(x)$. Around $x = 0$, p_0 behaves like $\frac{x^{\frac{2\lambda}{\sigma^2}}}{\gamma(x)^2} \exp\left(2\int_\cdot^x \frac{g(y)}{\gamma(y)^2}\,dy\right)$. The term in the exponential behaves like $\frac{\lambda}{\sigma^2} y^{\kappa}$. The integrability at zero hence depends on the exponent $-2(1 - \frac{\lambda}{\sigma^2})$: p_0 is integrable if the exponent is strictly larger than -1, i.e. $2\lambda/\sigma^2 > 1$. In the case $2\lambda/\sigma^2 < 1$, the non-integrability is due to a polynomial divergence, and hence might be compensated by a suitable choice of K. The only possible choice of K compensating the divergence at zero would correspond to:

$$p(1) = 2K \int_1^0 y^{-2\lambda/\sigma^2} \exp\left(-2\int_1^y \frac{g(z)}{\gamma(z)^2}\,dz\right) dy,$$

and here again the integral diverges at zero, and hence there is no such solution.

At infinity, the exact same analysis as done in the case $\alpha > 1$ applies, and hence we obtain that p_0 is integrable at ∞ under assumption (H6). $\square$

2.5. Heuristic discussion

We hence proved that as a function of the value of the coefficient α, the system can be in one of two substantially different regimes: in the case $\alpha < 1$ the singular point is always stable, and reached in finite time, and for $\alpha > 1$ the stability of the singular point is not affected by the presence of noise and only depends on the stability of the fixed point in the noiseless system.

The case $\alpha = 1$ constitutes the transition between these two regimes, and in that case the stability of the singular point depends both on the eigenvalue of the Jacobian matrix of the drift at the singular point and on the level of noise.

In order to understand heuristically these stabilization and destabilization phenomena, let us focus on the behavior of the system in the neighborhood of the singular point. In the case $\alpha > 1$, the diffusion coefficient vanishes faster than the drift coefficient at the singular point, and the system locally behaves as if there were no noise in the system. In particular, the stability of the singular point remains unchanged from that of the noiseless system. On the contrary, when $\alpha < 1$, the diffusion coefficient vanishes slower than the drift. When the system approaches the singular point, it is hence mainly driven by random fluctuations related to the diffusion coefficient. When the system is brought away from the singular point, noise increases, and when it is brought towards the singular point noise decreases. This induces a ratchet-like phenomenon: consider to points A and B close of the singular point, A being closer than B. The time it takes to go from A to B with power diffusion coefficient is way larger than the time needed to go from B to A. This behavior of the noise close of zero explains the stability in probability, and acts as a weak drift by 'stabilizing' the solutions when they approach the singular point. The intermediate case $\alpha = 1$ is precisely the transition where these two effects compensate. When the noise coefficient is large enough compared to the eigenvalue of the Jacobian matrix at the singular point, the noise effects described for $\alpha < 1$ dominate and stabilize the fixed point, and on the contrary the deterministic phenomena govern the dynamics similarly to the case $\alpha > 1$.

3. The stochastic pitchfork, Hopf and saddle-node bifurcations

Now that we characterized in detail the solutions of equation (1), we are in a position address the dynamics of three prominent bifurcations occurring in the study of dynamical systems: the pitchfork, Hopf and saddle-node (fold) bifurcation with power diffusion coefficients.

3.1. Analysis of the stochastic pitchfork bifurcation

In this section we first study the dynamics of an SDE with drift given by the normal form of the pitchfork bifurcation, before addressing the universality of these behaviors.

3.1.1. Dynamics

We consider the supercritical stochastic pitchfork bifurcation with power diffusion.

$$dx_t = (\lambda x_t - x_t^3)\, dt + \sigma |x_t|^\alpha dW_t \tag{10}$$

for $x_t \in \mathbb{R}$, with initial condition X_0 at $t = 0$. In this equation, λ is a real parameter, σ is a non-negative parameter and $\alpha \geq \frac{1}{2}$. In the case $\sigma = 0$, it is well known that the solution $x = 0$ is a stable fixed point for any $\lambda < 0$, and unstable for $\lambda > 0$, and two additional stable equilibria $\pm\sqrt{\lambda}$ exist in the region $\lambda > 0$ (see e.g. [1]).

For $\sigma \neq 0$, the model clearly satisfies assumptions (H1)-(H6) with $\nu = -\mu = 1 > 0$, $\kappa = \beta = 2$, $d = \sigma$ and $\delta = \alpha$. Moreover, it satisfies the additional assumptions used in proposition 4 since $g(x) = -x^3$ is differentiable with bounded derivative at zero and divergence at infinity upperbounded by x^4. All results of section 2 apply and we have:

Theorem 4.
- *For $\alpha < 1$, the fixed point δ_0 is stable in probability. Moreover, any solution almost surely reaches zero in finite time. Quasi-stationary distributions exists for $\alpha \in (1/2, 3/4)$.*

- *For* $\alpha = 1$ *and :*
  - $\lambda < \sigma^2/2$, δ_0 *is almost surely exponentially stable and no additional stationary solution exist.*
  - $\lambda \geq \sigma^2/2$, δ_0 *is unstable in probability, and two stationary solutions stable in the first approximation exist.*
- *for* $\alpha > 1$ *and:*
  - $\lambda < 0$, δ_0 *is stable in probability and no other stationary solution exist.*
  - $\lambda > 0$, δ_0 *unstable in probability and stationary solutions exist.*

Proof. Most results presented in the theorem are direct applications of the results demonstrated in section 2. The only assertion to be demonstrated is the stability in the first approximation of the additional stationary solutions.

In the case $\alpha = 1$, these solutions have the expression:

$$\mu_{\pm}(x) = \frac{2\sigma^{1-\frac{2\lambda}{\sigma^2}}}{\Gamma\left(-\frac{1}{2}+\frac{\lambda}{\sigma^2}\right)} x^{-2(1-\frac{\lambda}{\sigma^2})} e^{-x^2/\sigma^2} \mathbb{1}_{\pm x \geq 0}.$$

where we made explicit the normalization constant. The linearized equation reads:

$$dv_t = (\lambda - 3\,X_t^2)v_t\,dt + \sigma v_t\,dW_t \tag{11}$$

which is integrated as

$$v_t = \exp\left[(\lambda - \frac{\sigma^2}{2})t - \int_0^t 3\,X_s^2\,ds + \sigma W_t\right]v_0,$$

and hence the Lyapunov exponent is given by:

$$\limsup_{t\to\infty} \quad \frac{1}{t}\log|v_t| = (\lambda - \frac{\sigma^2}{2}) + \sigma\frac{W_t}{t} - \frac{1}{t}\int_0^t 3\,X_s^2\,ds + \frac{1}{t}\log(v_0).$$

Using the strong law of large numbers [17, Chap.IV.5], the term $\frac{1}{t}\int_0^t 3\,X_s^2\,ds$ converges a.s. towards $3\mathbb{E}[X_t^2]$ which can be computed in closed form, yielding the following expression of the Lyapunov exponent:

$$\begin{aligned} l &= (\lambda - \frac{\sigma^2}{2}) - 3\frac{\frac{1}{2}\sigma^{-1+\frac{2\lambda}{\sigma^2}}\Gamma\left(-\frac{1}{2}+\frac{\lambda}{\sigma^2}\right)}{\frac{1}{2}\sigma^{1+\frac{2\lambda}{\sigma^2}}\Gamma\left(\frac{1}{2}+\frac{\lambda}{\sigma^2}\right)} \\ &= (\lambda - \frac{\sigma^2}{2}) - 3\,(\lambda - \frac{\sigma^2}{2}) \\ &= -2\,(\lambda - \frac{\sigma^2}{2}) < 0 \end{aligned}$$

which ensures stability in the first approximation of the new stationary solutions since $\lambda > \frac{\sigma^2}{2}$. □



(a) $\alpha = 0.6$ (b) $\alpha = 0.85$

Figure 2: Distribution (ordinate truncated for legibility)of the trajectories that did not reach zero in the subcritical pitchfork model with $\lambda = -0.5$, $\sigma = 0.5$ different values of α, computed at 6 times. (a)$\alpha = 0.6$: stabilization on a quasi-stationary distribution that do not depend on time, $\alpha = 0.85$: the repartition varies as a function of time and all trajectories approach zero, arguing for the non-existence of quasi-stationary distribution. Simulations were done using a particle method with 100 000 particles, using the Euler-Maruyama scheme with $dt = 0.01$.

Let us now further describe the dynamics of the solutions as a function of α.

- **Case** $\alpha = 1$**:** We observe that for any $\lambda > 0$, the unstable deterministic fixed point becomes asymptotically exponentially stable when the noise parameter σ is large enough. For $\lambda > \frac{\sigma^2}{2}$, two symmetrical stationary distributions appear. For $\frac{\sigma^2}{2} < \lambda \leq \sigma^2$ the stationary distribution concentrates at zero and has a non increasing density diverging at zero. For $\lambda \geq \sigma^2$, the probability density function of the stationary distribution vanishes at zero and has a unique maximum reached for $x = \pm\sqrt{\lambda - \sigma^2}$ (see Figure 3). There is hence a qualitative transition at $\lambda = \sigma^2$, or P-bifurcation. In comparison with the deterministic bifurcation, the loss of stability is delayed and noise tends to stabilize the saddle point.

- **Case** $\alpha > 1$**:** the deterministic picture is qualitatively and quantitatively recovered: for $\lambda < 0$, δ_0 is stable in probability and is the unique stationary solution, and for $\lambda > 0$, δ_0 is unstable in probability, two additional stationary solutions appear, presumably stable as shown for $\alpha = 2$. The stationary probability density functions for $\lambda > 0$ vanish at zero and reach a maximum at X_m the solution of

  $$\left(\sigma^2\alpha - \lambda X^{2-2\alpha} + X^{-2\alpha+4}\right) = 0.$$

  The behavior of x_m when λ is close to zero follows one of the two regimes:

  - for $\alpha < 2$, $X_m \sim \left(\frac{\lambda}{\alpha\sigma^2}\right)^{1/(2\alpha-2)}$. For instance for $\alpha = \frac{3}{2}$ we get $X_m = -\frac{3}{4}\sigma^2 + \frac{1}{4}\sqrt{9\sigma^4 + 16\lambda}$
  - for $\alpha > 2$, $X_m \sim \sqrt{\lambda}$ and is insensitive to noise parameters σ and α. For $\alpha = 3$ we obtain the explicit form $X_m = \frac{1}{\sqrt{6}}\frac{\sqrt{-1+\sqrt{1+12\sigma^2\lambda}}}{\sigma}$
  - for $\alpha = 2$, this maximum can be computed explicitly, and is reached for $x = \sqrt{\frac{\lambda}{1+2\sigma^2}}$

  Surprisingly, in the limit $\lambda \to 0$ there is a discontinuity in the behavior of X_m as a function of α at $\alpha = 2$. Nevertheless, when $\lambda > 0$ this discontinuity is smoothed out. We conclude



Figure 3: (a) Codimension 2 stochastic bifurcation diagram of the pitchfork bifurcation: orange: zero is stable, blue: zero is unstable, yellow: zero is stable and is reached in finite time. (b) Sample path trajectories in all 6 regimes for the stochastic pitchfork bifurcation. Parameters $(\lambda, \alpha, \sigma)$: (b1) $(10, 0.6, 1)$ (quasi-stationary distribution plotted in Fig. 2(a)) (b2): $(2, 1, 1.2)$, (b3): $(1, 1.5, 1.2)$, (b4): $(0.5, 1, 1.2)$, (b5): $(-1, 1.5.1.2)$. In (b2) and (b3) the peak of the stationary distribution is depicted in dashed black. (c) Representation of stationary densities for the stochastic pitchfork system with linear noise, as a function of the parameter λ. Red square: δ_0 is stable. Blue and colored surfaces display the probability distribution $p_\lambda(x)$ of the stable stationary distributions, and the black line represents the maximal value of the distribution. The case $\alpha = 1$ was also studied using the theory of random dynamical systems in [9].

that in the case $\alpha > 2$, the behavior of the system is close from the one of the deterministic pitchfork bifurcation and the stationary distribution escapes zero as a square root of the parameter. But when $1 < \alpha < 2$, this is no more the case: the scaling of distribution peak behaves as $\lambda^{1/(2\alpha-2)}$, hence much slower than $\sqrt{\lambda}$. As σ is increased, the peak of the stationary distribution gets closer to zero.

- **Case** $\alpha < 1$**:** the solution δ_0 is always stable in probability. This result can appear relatively surprising at first sight. Indeed, in the deterministic case, λ is the exponential rate of divergence from the solution 0. However, adding a (possibly small) diffusion term proportional to $|x|^\alpha$ with $\alpha < 1$ stabilizes δ_0 in probability whatever the value of the noise intensity $\sigma \neq 0$. This observation, added to the fact that the solution is not exponentially asymptotically stable (though stable in probability) raises the question of how this convergence occurs. Figure 3(b1) presents a typical sample path of the process. Starting from a positive initial condition, we observe that the solution is evolving in the half-plane $x > 0$ and does not show any extinction clue. However, it suddenly reaches zero where it is absorbed after that random transient phase. This perfectly illustrates a typical behavior of the solutions of the pitchfork equation for $\alpha < 1$: the extinction time was shown to be almost surely finite. Moreover, trajectories that did not hit zero are distributed according to a quasi-stationary distributions as long as $\alpha < 3/4$.

Remark. *Let us briefly discuss the case of the subcritical stochastic pitchfork bifurcation:*

$$dx_t = (\lambda x_t + x_t^3)\,dt + \sigma |x_t|^\alpha dW_t.$$

The analysis of this system can be treated in the same fashion as the supercritical case. In the deterministic case ($\sigma = 0$*),* 0 *is stable for* $\lambda < 0$ *and unstable for* $\lambda > 0$*. Two additional unstable fixed points* $\pm\sqrt{-\lambda}$ *exist in the region* $\lambda < 0$*, and for* $\lambda > 0$ *the solution of the deterministic pitchfork equations blow up in finite time. In the case* $\alpha = 1$*, the analysis of the stochastic case gives the following conclusions. The fixed point* 0 *of the subcritical stochastic pitchfork equation is almost surely exponentially unstable for* $\lambda > \frac{\sigma^2}{2}$*, asymptotically unstable in probability for* $\lambda > -\frac{\sigma^2}{2}$ *and asymptotically stable in probability if* $\lambda < -\frac{\sigma^2}{2}$*. No extra stationary solution appear in that case.*

3.1.2. Universality properties

Similarly to the analysis of nonlinear dynamical systems, the analysis of the pitchfork equation reveals universal features of a wide class of systems. We formally show the universality property of the pitchfork bifurcation for general SDE on $\mathbb{R}$:

$$dx_t = f(x_t, \lambda)\,dt + g(x_t) dW_t$$

with a smooth f (at least three times continuously differentiable). We consider the dynamics of the system for parameter values λ in a neighborhood of $\lambda_0 \in \mathbb{R}$. Let us assume that:

- there exists $\nu > 0$ and $\kappa > 0$ such that for all λ in a neighborhood of λ_0, $f(x, \lambda) \leq -\nu x^{1+\kappa}$,
- for any λ, f is an odd function: $f(-x, \lambda) = -f(x, \lambda)$
- $\frac{\partial^3 f}{\partial x^3}(0, \lambda_0) \neq 0$

Defining $\xi_t = \beta(\lambda) x_t$ with $\beta(\lambda) = \sqrt{\frac{1}{6}\left|\frac{\partial^3 f}{\partial x^3}(0, \lambda)\right|}$, we have:

$$d\xi_t = \left(\frac{\partial f}{\partial x}(0, \lambda)\xi_t + \varepsilon \xi_t^3 + \Psi(\xi_t, \lambda)\right) dt + \beta(\lambda) g\left(\frac{\xi_t}{\beta(\lambda)}\right) dW_t.$$

where $\Psi(x,\lambda) = O(x^5)$ and $\varepsilon = \mathrm{sign}(\frac{\partial^3 f}{\partial x^3}(0,\lambda))$ which is well defined around $\lambda = \lambda_0$.

Let us now assume that $g(x) \sim \sigma|x|^\alpha$ around $x = 0$ and $g(x) > 0$ for $x \neq 0$. We can hence apply the results developed in the previous section to characterize the dynamics of this general system. Pitchfork-like transitions occur at $\lambda = \lambda_0$ when:

- $\alpha > 1$, $\frac{\partial f}{\partial x}(0,\lambda_0) = 0$ and $\frac{\partial^2 f}{\partial x \partial \lambda}(0,\lambda_0) \neq 0$ or
- $\alpha = 1$, $\frac{\partial f}{\partial x}(0,\lambda_0) = \frac{\sigma^2}{2}$ and $\frac{\partial^2 f}{\partial x \partial \lambda}(0,\lambda_0) \neq 0$.

We now put in good use this reduction in the following example.

Example 1. *Let us consider the Wilson and Cowan firing-rate model [19], classically describing the behavior of a cortical column. In that model, the voltage v of a typical neuron in the column is solution of the equation:*

$$dv_t = \left(-\frac{1}{\tau} v_t + J S(v_t)\right) dt + \gamma(v_t) dW_t$$

where $S(x)$ is a smooth sigmoidal function describing the voltage-to-rate transformation, and J is the typical connectivity strength. We choose here $S(x) = \tanh(g\,x)$ with $g > 0$ the sharpness coefficient of the sigmoid and $\gamma(x) = \sigma|x|^\alpha$ with $\alpha \geq \frac{1}{2}$. This equation can be locally reduced to the normal form of the pitchfork bifurcation since the flow is symmetrical, and we obtain using the same transformation as performed above:

$$d\xi_t = \left((-\frac{1}{\tau} + gJ)\xi_t - \xi_t^3 + \Psi(\xi_t, J, g)\right) dt + 2^{1-\alpha}\sigma|\xi_t|^\alpha dW_t$$

For $\alpha < 1$, the fixed point 0 is almost surely reached in finite time, for $\alpha > 1$, the fixed point 0 is stable in probability for $gJ\tau < 1$ and unstable in probability otherwise. For $\alpha = 1$, the fixed point is stable in probability if $-1/\tau + gJ < \sigma^2/2$ and unstable otherwise. This behavior is illustrated numerically in Figure 4.

3.2. The stochastic Hopf bifurcation

The Hopf bifurcation for smooth dynamical system is topologically equivalent to the two dimensional normal form:

$$\frac{\mathrm{d}x}{\mathrm{d}t} = \beta\, x - y + \varepsilon x(x^2 + y^2) \qquad ; \qquad \frac{\mathrm{d}y}{\mathrm{d}t} = \beta\, y + x + \varepsilon y(x^2 + y^2)$$

where β is the bifurcation parameter and $\varepsilon = \pm 1$ governs the type of bifurcation. It can be conveniently written in complex form for $z_t = x_t + \mathbf{i}\, y_t$ with $\mathbf{i}^2 = -1$:

$$\frac{\mathrm{d}z}{\mathrm{d}t} = (\beta + \mathbf{i} + \varepsilon|z|^2)\, z.$$

The null solution ($x = 0, y = 0$) is always solution of the equations. It is exponentially stable for any $\beta < 0$ and exponentially unstable for any $\beta > 0$. If $\varepsilon = -1$, the system presents stable



(a) $\alpha = 0.8$ (b) $\alpha = 1$ (c) $\alpha = 1.5$

Figure 4: Wilson and Cowan system with power diffusion function: empirical mean m of v_T at time $T = 200$ for 1 000 independent sample path as a function of g. $\alpha < 1$: the fixed point 0 is reached in finite time whatever g, $\alpha = 1$: zero looses stability for $g = 1.5$ and $\alpha > 1$: zero looses stability for $g = 1$. Due to the finite simulation time, the transitions appear smooth in these simulations. Euler Maruyama scheme with time step $dt = 0.001$, 100 values for parameter g.

oscillations for $\beta > 0$ and the bifurcation is termed supercritical, and if $\varepsilon = 1$, unstable periodic orbit exist for $\beta < 0$ and the bifurcation is termed subcritical.

We study in this section the stochastic Hopf bifurcation with multiplicative noise. For simplicity, we will consider that the stochastic perturbations are driven by a single Brownian motion W_t. Let $\sigma^* = \sigma + \mathbf{i}\mu$ be a noise parameter, the stochastic Hopf normal form equation reads (in complex form):

$$dZ_t = (\beta + i + \varepsilon |Z_t|^2) Z_t\, dt + \sigma^* |Z_t|^{\alpha - 1} Z_t\, dW_t,$$

which corresponds for $Z_t = X_t + \mathbf{i} Y_t$ to the bidimensional system:

$$\begin{cases} dX_t &= (\beta X_t - Y_t + \varepsilon X_t\, (X_t^2 + Y_t^2))\, dt + (\sigma X_t - \mu Y_t) |Z_t|^{\alpha-1}\, dW_t \\ dY_t &= (\beta Y_t + X_t + \varepsilon Y_t\, (X_t^2 + Y_t^2))\, dt + (\mu X_t + \sigma Y_t) |Z_t|^{\alpha-1}\, dW_t \end{cases} \tag{12}$$

where β and ε correspond to the parameters of the Hopf bifurcation and μ and σ are two parameters globally governing the amplitude of the stochastic perturbation.

The deterministic process $Z_t = 0$ corresponding to $X_t = 0, Y_t = 0$ for all $t \geq 0$ is solution of (12) and is univocally defined by the fact that the modulus of Z_t is null. We denote by R_t the modulus of Z_t and by θ_t its argument[2].

Lemma 1. *The modulus of the variable $R_t = |Z_t|$ and the argument θ_t satisfy the equations:*

$$\begin{cases} dR_t &= \left(\beta R_t + \frac{\mu^2}{2} R_t^{2\alpha-1} + \varepsilon R_t^3\right) dt + \sigma R_t^{\alpha}\, dW_t \\ d\theta_t &= (-\sigma\mu R_t^{2\alpha-2} + 1)\, dt + \mu R_t^{\alpha-1} dW_t \end{cases}$$

Proof. Let (X_t, Y_t) be a solution of the Hopf equations (12). We apply Itô formula to the variable

---

[2] $R_t = |Z_t| = \sqrt{X_t^2 + Y_t^2}$, θ_t given by the $\arctan(\frac{Y_t}{X_t})$ when $X_t > 0$, added or subtracted π when $X_t < 0$ depending on the sign of Y_t, and $\pm\pi/2$ if $X_t = 0, \pm Y_t > 0$

$R_t = \sqrt{X_t^2 + Y_t^2}$:

$$
\begin{aligned}
dR_t &= \frac{1}{R_t}(X_t dX_t + Y_t dY_t) + \frac{1}{2\,R_t^3}\left(Y_t^2 d\langle X\rangle_t + X_t^2 d\langle Y\rangle_t + 2\,X_t\,Y_t d\langle X, Y\rangle_t\right) \\
&= \Big\{\frac{1}{R_t}\Big(\beta\,X_t^2 - X_t\,Y_t + \varepsilon X_t^2\,(X_t^2 + Y_t^2) + \beta\,Y_t^2 + X_t\,Y_t + \varepsilon Y_t^2\,(X_t^2 + Y_t^2)\Big) \\
&\quad + \frac{R_t^{2\alpha-2}}{2\,R_t^3}\Big(Y_t^2(\sigma\,X_t - \mu\,Y_t)^2 + X_t^2(\sigma\,Y_t + \mu\,X_t)^2 - 2X_t\,Y_t\,(\sigma\,X_t - \mu\,Y_t)\,(\sigma\,Y_t + \mu\,X_t)\Big)\Big\}\,dt \\
&\quad + \frac{R_t^{\alpha-1}}{R_t}\Big\{X_t\,(\sigma\,X_t - \mu\,Y_t) + Y_t\,(\sigma\,Y_t + \mu\,X_t)\Big\}\,dW_t \\
&= \Big\{\frac{1}{R_t}(\beta R_t^2 + \varepsilon R_t^4) + \frac{1}{2R_t^3}\mu^2 R_t^{4+2\alpha-2}\Big\}\,dt + \frac{1}{R_t}\sigma\,R_t^{2+\alpha-1}\,dW_t \\
&= \left(\beta R_t + \frac{\mu^2}{2}R_t^{2\alpha-1} + \varepsilon R_t^3\right)\,dt + \sigma\,R_t^{\alpha}\,dW_t
\end{aligned}
$$

The argument θ_t is given by $\theta_t = \arctan(Y_t/X_t)$ plus possibly constants depending on the sign of X_t and Y_t. Applying Itô's formula again yields:

$$
d\theta_t = (1 - \sigma\mu R_t^{2(\alpha-1)})\,dt + \mu R_t^{\alpha-1} dW_t
$$

which ends the proof of the lemma. □

It is important to note that the equation on the modulus is uncoupled of the phase equation on θ_t. The modulus of (X_t, Y_t) is therefore solution of a stochastic differential equation of type bifurcation (1). Moreover, when $\alpha = 1$, the equations take the simpler form:

$$
\begin{cases}
dR_t &= \left((\beta + \frac{\mu^2}{2})R_t + \varepsilon R_t^3\right)\,dt + \sigma\,R_t\,dW_t \\
d\theta_t &= (1 - \sigma\mu)\,dt + \mu dW_t
\end{cases}
$$

and hence the variable R_t is solution of a stochastic pitchfork bifurcation. By application of theorems 4 and remark 3.1.1, it is easy to establish the following:

Theorem 5. *For $\alpha = 1$, the null solution of the supercritical Hopf equations is almost surely exponentially stable if $\beta < \frac{\sigma^2-\mu^2}{2}$ and asymptotically stochastically unstable if $\beta > \frac{\sigma^2-\mu^2}{2}$. In that case, there exists a new stochastically stable stationary solution with distribution:*

$$
p_R(x) = \frac{2\sigma^{1-\frac{2\lambda}{\sigma^2}}}{\Gamma\left(-\frac{1}{2} + \frac{\lambda}{\sigma^2}\right)} x^{-2(1-\frac{\lambda}{\sigma^2})} e^{-x^2/\sigma^2}\,\mathbb{1}_{x\geq 0}.
$$

The null solution of the subcritical Hopf equations is almost surely exponentially unstable if $\beta > \frac{\sigma^2-\mu^2}{2}$. It is asymptotically stochastically unstable if $\beta > -\frac{\sigma^2+\mu^2}{2}$ and stochastically stable if $\beta < -\frac{\sigma^2+\mu^2}{2}$.

For $\sigma \neq 0$, the modulus R_t is a.s. exponentially attracted towards 0 when $\beta < \frac{\sigma^2-\mu^2}{2}$. Therefore, as the imaginary part of the noise, σ, is increased, the fixed point gains stability. For



Figure 5: Simulations of sample path trajectories of (X_t, Y_t) for the stochastic Hopf bifurcation with $\alpha = 1$, $\mu = 0$ and $\sigma = 1$ for different values of β (deterministic phase case). (A):$\beta = 0.3 < \frac{\sigma^2}{2}$: zero is a.s. exponentially stable; (A1): trajectories (blue: X_t, red: Y_t), (A2): phase plane, (A3): phase θ_t is deterministic equal to t: the solutions rotate with pulsation one. This is true of all three cases (A), (B) and (C). (B):$\beta = 0.7 \in [\frac{\sigma^2}{2}, \sigma^2]$: Zero is unstable and the stationary modulus distribution is peaked at zero; (B1): trajectories, (B2): phase plane, (B3): distribution of the modulus R_t. (C): $\beta = 4 > \sigma^2$: zero is unstable and the distribution of the modulus has a peak around $R^* > 0$ (C3): the trajectories display perfectly periodic oscillations (C1, C2) with random modulus, a highly regular behavior for a stochastic system.

$\beta \in (\frac{\sigma^2-\mu^2}{2}, \frac{2\sigma^2-\mu^2}{2})$, the fixed point zero is unstable and solutions do not converge towards the fixed point zero. The distribution of the moduli R_t converge towards the stationary distribution $p_R(x)$ heavily charging zero, and solutions stay close from the fixed point with random stochastic variations. As soon as $\beta > \frac{2\sigma^2-\mu^2}{2}$, the solutions have their modulus centered at $R^* := \sqrt{\beta - \frac{2\sigma^2-\mu^2}{2}}$ and present stochastic oscillations with a deterministic pulsation $(-1 + \sigma\mu)$

and random phase given by a Brownian motion scaled by μ. When $\mu = 0$, the phase is hence deterministic and the system rotates with pulsation 1. When $\beta \in (\frac{2\sigma^2-\mu^2}{2}, \frac{\sigma^2-\mu^2}{2})$, the stationary modulus of the solution has a distribution peaked at zero and no actual oscillation can be observed. When $\beta > \frac{2\sigma^2-\mu^2}{2}$, the stationary modulus of the solution has a maximum at $R^* = \sqrt{\beta - \frac{2\sigma^2-\mu^2}{2}}$ and the distribution is null at zero: solutions hence correspond to perfectly periodic oscillations with random amplitude. Interestingly, when $\sigma\mu$ crosses 1, the direction of the oscillation changes: for $\sigma\mu < 1$, the oscillation is counter clockwise, and for $\sigma\mu > 0$, the oscillation is clockwise. This is a new type of bifurcation due to noise that we term orientation bifurcation. This phenomenon is illustrated in Figure 6. In the case $\sigma\mu = 1$, the pulsation is null and the phase is given by μW_t: it has a probability $1/2$ to rotate clockwise and $1/2$ to rotate counterclockwise, and changes the sense of rotation. The rotation number is equal to zero.



(a) $\sigma = 0.5$

(b) $\sigma = 1$

(c) $\sigma = 1.5$

Figure 6: Noise-induced inversion of the rotation pulsation: $\alpha = 1, \beta = 4, \mu = 1$ for different values of σ. Blue: phase (θ_t) as a function of time and black: deterministic pulsation $1 - \sigma\mu$. (a) counter-clockwise rotation, (b): no rotation, trajectories appear stochastic and the phase is a Brownian motion, and (c): clockwise rotation.

We observe in the purely imaginary noise case ($\sigma = 0$), R_t satisfies the deterministic pitchfork bifurcation equation. For $\beta > \frac{\mu^2}{2}$, $R_t = \sqrt{\beta + \frac{\mu^2}{2}}$ is the only attractive solution: the solution of the stochastic Hopf bifurcation asymptotically lives on a deterministic circle. Its rotates on this circle with a stochastic phase given by a Brownian motion multiplied by μ and a deterministic pulsation equal to 1 (see Figure 7).

Let us eventually note the opposite effects of the real and imaginary values of the noise parameter: for fixed β, increasing σ tends to stabilize 0, whereas increasing μ has the effect of destabilizing 0. These two effects compensate each other, and when choosing $\sigma = \mu$, the loss of stability of zero arises at $\beta = 0$ precisely as it is the case of the deterministic bifurcation. However, cycles appear for $\beta > \sigma^2/2$ in that case, and hence we observe a systematic delay in the apparition of cycles for $\sigma \neq 0$.

Beyond these interesting effects of noise, we want to underline the fact that for $\alpha = 1$ the global picture of the deterministic Hopf bifurcation is nevertheless conserved, since as β is increased, a limit cycle appears, around which the solutions stochastically rotate. The picture will be relatively different in the nonlinear noise case $\alpha \neq 1$. In that case we prove the following:

Proposition 6. *In the case of the supercritical ($\varepsilon = -1$) stochastic Hopf bifurcation, we have:*

- *For $\alpha < 1$, the fixed point 0 is:*
  - *stable in probability for $|\mu| < |\sigma|$*
  - *unstable in probability for $|\mu| > |\sigma|$*



Figure 7: Simulations of sample path trajectories of (X_t, Y_t) for the stochastic Hopf bifurcation with $\alpha = 1$, $\sigma = 0$ and $\mu = 2$ (deterministic modulus case) for different values of β. (A):$\beta = -2 < -\frac{\mu^2}{2}$: zero is a.s. exponentially stable; (A1): trajectories (blue: X_t, red: Y_t, black: $R_t s$), (A2): phase plane, (A3): argument θ_t in blue, deterministic drift in black. (B): $\beta = 0 > -\frac{\mu^2}{2}$: the modulus converges to a constant value $\sqrt{\mu^2/2} = \sqrt{2}$; (B1): trajectories, (B2): phase plane, (B3): Argument θ_t (blue), deterministic drift (black).

Moreover, almost all sample path reaches zero in finite time.

- *For* $\alpha > 1$*, the fixed point* 0 *is:*
  - *stable in probability for* $\beta < -\frac{\mu^2}{2}$
  - *unstable in probability otherwise. Two additional stationary solutions exist in that case.*

The solutions almost surely never reach zero.

Remark. *Let us emphasize the fact that the case* $\alpha < 1$ *is substantially different from the general case: indeed, we had seen that the singular point was always stochastically stable. Surprisingly here the stability of zero depends on the relative values of the real and imaginary parts of the noise.*

Proof. The case $\alpha > 1$ falls in the general analysis developed in section 2. Indeed, the diffusion is of the form of equation (1) with $g(x) = \frac{\mu^2}{2} x^{2\alpha - 1} - x^3$ which clearly satisfies assumptions (H1) and (H2). The general analysis hence applies and directly leads to the conclusion of the proposition.

In the case $\alpha < 1$, the equation was not covered in the general analysis, since the drift includes a term $\frac{\mu^2}{2} x^{2\alpha-1}$ and $2\alpha - 1 < 1$. However, the same principles may be applied to that case. First of all, existence and uniqueness of solutions is proved using the same arguments, and the a.s. boundedness of the process as well. Hence there exist a unique solution to the stochastic Hopf bifurcation process with $\alpha < 1$, which is almost surely bounded for all times. The stability of zero is investigated as follows. For $\mu > \sigma$, the Lyapunov function $V(x) = -\log(x)$ is C_0^2 around zero, positive, diverges at zero, and

$$\mathcal{L}V(x) = \left(-\frac{\mu^2}{2} + \frac{\sigma^2}{2}\right) x^{2\alpha-2} + o(x^{2\alpha-2})$$

and is hence negative around zero, proving that zero is unstable in probability for $\mu > \sigma$.

For $\mu < \sigma$, we define the Lyapunov function $V(x) = x^{\Phi}$ with $\Phi = \frac{\sigma^2-\mu^2}{2\sigma^2} > 0$. This function is C_0^2 and we have:

$$\begin{aligned}\mathcal{L}V(x) &= \Phi\left(\frac{\mu^2}{2} + (\Phi - 1)\frac{\sigma^2}{2}\right) x^{2\alpha-2+\Phi} + o(x^{2\alpha-2+\Phi})\\ &= \Phi\frac{\mu^2 - \sigma^2}{4} x^{2\alpha-2+\Phi} + o(x^{2\alpha-2+\Phi})\end{aligned}$$

and hence is negative close of zero. Using the same techniques as in the general theory, it is easy to conclude that the singular point is unstable in probability. In order to prove that the solutions almost surely hit zero in finite time, we proceed as in the proof of proposition 2. We know that the process is almost surely bounded, and analyze Feller's scale functions. It is clear that $p(x) \to \infty$ as $x \to \infty$ following the proof of proposition 2: the proof only uses the higher order term $-x^3$. We hence need to ensure that $\lim_{x\to 0^+} v(x) < \infty$. This property is related to fine properties of the scale functions close from zero, which is now governed by the term $\mu^2/2\ x^{2\alpha-1}$. We hence have, around 0:

$$G(x) = \frac{\mu^2}{2\sigma^2}\log(x) + \frac{\lambda}{\sigma}\frac{x^{2-2\alpha}}{2(1-\alpha)} - \frac{x^{2(2-\alpha)}}{2(2-\alpha)\sigma^2}$$

and therefore we can obtain the explicit expression of $p(x)$:

$$p(x) = \int_c^x y^{-\frac{\mu^2}{\sigma^2}} \exp\left(-\frac{\lambda}{\sigma^2(1-\alpha)} y^{2-2\alpha} + \frac{2y^{4-2\alpha}}{(4-2\alpha)\sigma^2}\right) dy$$

More importantly, in order to characterize v, we consider the term $G(y)-G(z)$ which is equivalent close from 0 to: $G(y) - G(z) \sim_{0^+} \frac{\mu^2}{2\sigma^2}\log(\frac{y}{z})$. This implies that:

$$\begin{aligned}\int_y^x \exp(2G(y) - G(z))\, dz &\sim_{0^+} \int_y^x \exp\left(\frac{\mu^2}{\sigma^2}\log(\frac{y}{z})\right) dz\\ &\sim \frac{1}{1-\mu^2/\sigma^2}\left(\left(\frac{y}{x}\right)^{\mu^2/\sigma^2} - 1\right)\end{aligned}$$

and hence:

$$v(x) \sim \frac{2}{\sigma^2}\left(\frac{x^2}{\mu^2/\sigma^2 + 1} - x\right)$$

This function will hence remain bounded as $x \to \infty$, ensuring using Feller's test for explosion that the first exit time of the interval $(0, \infty)$ is almost surely bounded. The almost sure boundedness of the process allows concluding that the probability first hitting time of zero is almost surely finite, which ends the proof. ☐

Therefore, similarly to the case of the stochastic pitchfork bifurcation, we observed that for $\alpha > 1$, the noise does not affect the stability of 0 and the loss of stability precisely appears for $\beta = 0$ as in the deterministic case. The behavior of the system beyond the bifurcation point depends on the shape of the stationary distribution, and in that case again the solution reaches a maximum for a positive value, reproducing qualitatively the behavior of the deterministic pitchfork bifurcation.

For $\alpha < 1$, the almost sure absorption at zero of the solution persists. However, the stochastic stability of the fixed point zero now depends on the relative values of μ and σ. This distinction is not fundamental as long as we are concerned with sample path properties, and similarly to the general case, almost any sample path will eventually be absorbed at zero in finite time.

Let us eventually remark that the universality properties of this normal form are harder to obtain than in the case of the pitchfork case already treated and the saddle-node bifurcation to come. This is not only due to the fact that we are in higher dimensions, but also (and this difficulty is probably harder to overcome) because the generic reduction to normal form in the Hopf bifurcation involves a nonlinear change of variable, which is not the case of the pitchfork or saddle-node bifurcation (see [2]). Since we consider diffusion processes, the process obtained through nonlinear change of variable will satisfy, using Itô formula, a SDE where the drift function involves a term related to the diffusion, and it is hence much more complex to obtain a suitable change of variable. Another approach would consist in showing that some function of the process satisfies an SDE of the type studied in section 2. Let us formally give the kind of computation that would be involved, considering a two-dimensional system:

$$dX_t = F(X_t)\, dt + G(X_t) dW_t$$

where $G(x)$ is precisely of the complex noise form considered above. The modulus of the solution R_t satisfies the SDE:

$$dR_t = \frac{1}{R_t}\left(X_t^1 F^1(X_t) + X_t^2 F^2(X_t) + \frac{\mu^2}{2} R_t\right) dt + (X_t^1 G^1(X_t) + X_t^2 G^2(X_t))dW_t$$

where $X^{1(2)}, F^{1(2)}$ and $G^{1(2)}$ represent the first (second) component of the vector field. The question that now arises is whether this system can be reduced to a pitchfork equation with possibly higher order terms, which can be checked in a case-by-case basis, as we now do for our choice example.

Example 2. *Let us consider a Wilson and Cowan neural network composed of an excitatory and an inhibitory population. The network equations in that case read:*

$$\begin{cases} dv_t^1 = (-v_t^1 + S(gv_t^1) - S(gv_t^2)) + (\sigma v_t^1 - \mu v_t^2)|v_t|^{\alpha-1} dW_t \\ dv_t^2 = (-v_t^2 + S(gv_t^1) + S(gv_t^2)) + (\sigma v_t^2 + \mu v_t^1)|v_t|^{\alpha-1} dW_t \end{cases} \tag{13}$$

with $S(x) = \tanh(g\, x)$ *and* $|v_t| = \sqrt{(v_t^1)^2 + (v_T^2)^2}$. *In the deterministic case, it is straightforward to show that the system undergoes a Hopf bifurcation at* $g = 1$. *Let us now consider the equation satisfied by the modulus* $\rho_t = |v_t|$. *Using Itô's formula we obtain:*

$$d\rho_t = \Big((-1 + g)\rho_t + \mu^2/2\rho_t^{2\alpha-1} + g(v_t^1, v_t^2)\Big)\, dt + \sigma \rho_t^{\alpha} dW_t$$

with

$$\begin{aligned} g(x,y) &= -\frac{g^3}{3\rho}(x^4 + y^4 + x^3y + xy^3) + O(\rho^4) \\ &= -\frac{g^3}{3\rho}((x+y)^4 - 2x^2y^2) + O(\rho^4) \leq -\frac{g^3}{12}\rho^3 + O(\rho^4) \end{aligned}$$

The argument satisfies the equation:

$$d\theta_t = (g - \sigma\mu\rho^{2\alpha-2} + O(\rho_t^2))\, dt + \mu\rho^{\alpha-1} dW_t$$

The dynamics of the system can hence be analyzed using the above analysis:

- $\alpha = 1$, *zero is stable in probability when* $(-1 + g + \mu^2/2) \leq \sigma^2/2$ *and unstable otherwise. At this point, a stochastic Hopf bifurcation arises, and oscillations appear, with pulsation locally equivalent to* $g - \mu\sigma$ *(with order two corrections in* ρ*) and stochastic phase given by a Brownian motion multiplied by* μ*. The oscillation orientation bifurcation takes place for* $g = \mu\sigma$*.*
- *For* $\alpha > 1$ *zero is stable if and only if* $(-1 + g + \mu^2/2) \leq 0$.
- *For* $\alpha < 1$, *the fixed point is stable in probability if* $|\mu| < |\sigma|$ *and unstable otherwise. The first hitting time of the singular point is almost surely finite in both cases.*

The rotation orientation transition is illustrated in Figure 8.

3.3. *The stochastic saddle-node bifurcation*

The third prominent universal codimension one bifurcation of equilibria in deterministic ODEs is the *saddle-node (or fold) bifurcation*. It corresponds to the normal form $\dot{x} = -x^2 + a$ (sometimes considered $x^2 - a$ which is equivalent to the normal form considered here through a change of time). This dynamical system has no equilibrium when $a < 0$ (solutions blows up in finite time) and two equilibria for $a > 0$: $\sqrt{a}$ which is stable and $-\sqrt{a}$ which is unstable. The system has multiple singular points, and these depends on the bifurcation parameter. Choices of diffusion functions vanishing only at $\sqrt{a}$ or at $-\sqrt{a}$ enter the application domain of the general theory developed in section 2. Let us consider now a case, contrasting with the previous cases treated, where the diffusion function vanishes at both singular points:

$$dX_t = (-X_t^2 + a)\, dt + \sigma|X_t^2 - a|^\alpha\, dW_t \tag{14}$$

For $a < 0$, there are two stationary distributions corresponding to the equilibria $\sqrt{a}$ and $-\sqrt{a}$. Denoting $y_t^+ = x_t - \sqrt{a}$ (vanishing at $x = \sqrt{a}$) and $y_t^- = x_t + \sqrt{a}$, it is easy to show that these variable satisfy the equations:

$$\begin{cases} dy_t^+ = (-2\sqrt{\alpha}y_t^+ - (y_t^+)^2)\, dt + \sigma|-2\sqrt{\alpha}y_t^+ - (y_t^+)^2|^\alpha\, dW_t \\ dy_t^- = (2\sqrt{\alpha}y_t^- - (y_t^+)^2)\, dt + \sigma|2\sqrt{\alpha}y_t^+ - (y_t^+)^2|^\alpha\, dW_t \end{cases}$$

In both case, the drift and diffusion coefficients satisfy the assumptions (H1), (H2), (H3), (H4) and (H6), and obviously assumption (H5) is not valid. Assumptions (H1), (H2) and (H6) are trivial since the function corresponding to $g(x)$ in the general cases is equal to $-x^2$, hence $\mu =$



(a) $\alpha = 0.8, \mu = 2, \sigma = 1$ (b) $\alpha = 1, \mu = 2\ \sigma = 1$ (c) $\alpha = 1, \mu = 1\ \sigma = 0.5$



(d) $\alpha = 0.8, \mu = 1, \sigma = 2$ (e) $\alpha = 1, \mu = 2\ \sigma = 1$ (f) $\alpha = 1, \mu = 1\ \sigma = 0.5$

Figure 8: Dynamics of the Wilson and Cowan two-populations system (13). (a)-(d): v^1_t (blue) and v^2_t (red) as a function of time; (e) and (f): θ_t (green) and deterministic drift (black) as a function of time. (a) and (d) correspond to $\alpha < 1$: for $|\mu| < |\sigma|$, the singular point is stable (d) and unstable otherwise (a). The other figures correspond to $\alpha = 1$ and illustrate the change of rotation orientation bifurcation at $\sigma\mu = g$. All the figures correspond to $g = 1$.

$-\nu = -1 < 0$ and $\beta = \kappa = 1$. The diffusion coefficients denoted $\gamma(x)$ in the general case satisfy for both the equation on y^+ and on y^-: $\gamma(x) \sim_{x\to 0^+} \sigma(2\sqrt{a})^\alpha x^\alpha$, ensuring that assumption (H3) is satisfied, and $\gamma(x) \sim_{x\to\infty} \sigma|y|^{2\alpha}$ ensuring that assumption (H4) is satisfied with $d = \sigma$ and $\delta = 2\alpha$.

In the deterministic case, for $a > 0$ and an initial condition smaller than $-\sqrt{a}$ or for any initial condition when $a < 0$, the solutions blow up in finite time to $-\infty$, and in usual applications additional, higher order confining terms prevent the blow up. Interestingly, in the present case, the presence of noise can prevent blow up as we now show:

Proposition 7. *The solutions of the stochastic saddle-node equation* (14) *are defined for all times or blow up in finite time under the conditions:*

- *For* $\frac{1}{2} \leq \alpha < 1$ *we have:*
  - *if* $a < 0$ *the solutions almost surely blow up in finite time to* $-\infty$
  - *if* $a > 0$, *the solutions with initial condition smaller than* $-\sqrt{a}$ *almost surely exit the interval* $(-\infty, -\sqrt{a})$ *in finite time.*
- *for* $\alpha \geq 1$, *the solutions almost surely never blow up, and for* $a > 0$, *the exit time of* $(-\infty, -\sqrt{a})$ *is almost surely infinite.*

Proof. We demonstrate this proposition using Feller's test for explosion.

Let us start by considering $a < 0$. The diffusion is then defined on $\mathbb{R}$ with no singular point, and Feller's scale function reads:

$$p(x) = \int_c^x \exp\left(-2 \int_c^y -(\xi^2 + a)^{1-2\alpha}\, d\xi\right) dy.$$

At infinity, the integrand inside the exponential is equivalent at infinity to $\frac{2}{2(2\alpha-1)-1} y^{-2(2\alpha-1)+1}$. This quantity has a finite limit when $\alpha \geq 1$, implying that $p(x) \to \pm\infty$ when $x \to \pm\infty$. Feller's test implies that the first exit time of $(-\infty, \infty)$ is infinite with probability one.

For $\alpha < 1$, $p(x)$ tends to ∞ at ∞, and has a finite limit at $-\infty$. This property ensures that the probability of the process diverge towards $-\infty$ is equal to one. It remains to prove that the explosion occurs in finite time. To this purpose, we need to show that the limit of Feller's scale function $v(x)$ is finite at $-\infty$. This is the case under the assumption $\alpha > 1$. Indeed, $v(x)$ is defined as:

$$v(x) = \int_c^x \int_c^y \exp\left(2 \int_z^y (u^2 - a)^{1-2\alpha}\right)$$

The integral in the argument of the exponential function is equivalent at $-\infty$ to $\frac{2}{\Phi}(y^{\Phi} - z^{\Phi})$ which tends to $-\infty$ as a power function, hence the exponential has moments of any order, and in particular is twice integrable at $-\infty$.

For $a > 0$, we consider the intervals $(-\infty, -\sqrt{a})$ and $(\sqrt{a}, \infty)$. The above analysis proved that for $\alpha \geq 1$, $p(x)$ tends to ∞ when $x \to \infty$ and towards a finite value at $x \to -\infty$. At $\pm\sqrt{a}$, the argument of the exponential term is equivalent to $\int x^{-2\alpha+1}$ which diverges for $\alpha \geq 1$, implying that the exit time of $(-\infty, -\sqrt{a})$, $(-\sqrt{a}, \sqrt{a})$ and $(\sqrt{a}, \infty)$ are all almost surely infinite and that there is no blow up in finite time. For $\alpha < 1$, both $p(-\sqrt{a}^-)$ and $p(\sqrt{a}^+)$ are finite. For $c \in (-\sqrt{a}, \sqrt{a})$, we have:

$$\begin{cases} G(x) = \int_c^x -(\xi + \sqrt{a})^{1-2\alpha}(-\xi + \sqrt{a})^{1-2\alpha}\, d\xi \\ v(x) = 2 \displaystyle\int_c^x \int_y^x \frac{\exp(-2(G(y) - G(z)))}{(-x^2 + a)^{2\alpha}}\, dz\, dy \end{cases}$$

and using a similar argument as used in the proof of proposition 2, we show that $\lim_{x \to \sqrt{a}^-} v(x) < \infty$, and by symmetry we hence have $\lim_{x \to -\sqrt{a}^+} v(x) < \infty$. Feller's test for explosion hence ensures that the first exit time of the interval $(-\sqrt{a}, \sqrt{a})$ is almost surely finite. We hence converge in finite time towards $\sqrt{a}$ or $-\sqrt{a}$. Moreover, the probability for a trajectory starting with an initial condition $x \in (\sqrt{a})$ to reach $\sqrt{a}$ (resp. $-\sqrt{a}$) prior to reaching $-\sqrt{a}$ (resp. $\sqrt{a}$), denoted $p_+(x)$ (resp. $p_-(x)$) have the expression:

$$p_-(x) = 1 - p_+(x) = \frac{p(\sqrt{a}^-) - p(x)}{p(\sqrt{a}^-) - p(-\sqrt{a}^+)}$$

This quantity is plotted in Figure 9(c). □

Application of the general results of section 2 yield the following:

Proposition 8. *For $a > 0$, the saddle-node equation* (14) *has two Dirac delta distributed functions $\delta_{\sqrt{a}}$ and $\delta_{-\sqrt{a}}$, which enjoy the stability properties:*

- *for $\alpha < 1$, both $\delta_{\sqrt{a}}$ and $\delta_{-\sqrt{a}}$ are stable in probability, and the first hitting time τ of one of the equilibria is almost surely finite.*

- *for* $\alpha = 1$, *the stationary distribution* $\delta_{\sqrt{a}}$ *is always stable in probability. The stationary distribution* $\delta_{-\sqrt{a}}$ *is stable in probability for* $a > \sigma^{-4}$ *and unstable in probability otherwise.*
- *for* $\alpha > 1$, *the stationary distribution* $\delta_{\sqrt{a}}$ *is stable in probability and* $\delta_{-\sqrt{a}}$ *is unstable in probability.*

Proof. This is a direct application of the results of section 2 applied to y^+ to characterize the properties of $\sqrt{a}$ and to y^- for $-\sqrt{a}$. For instance for the fixed point $-\sqrt{a}$ and $\alpha = 1$, the fixed point is stable if and only if $2\sqrt{a} < 2\sigma^2\alpha$ i.e. $\sigma^2\sqrt{a} > 1$. The finite absorption time property is a consequence of proposition 7. ☐

For $\alpha < 1$, we know that the trajectories either reach one of the equilibria or blow up in finite time, which describe perfectly the permanent regime of the saddle-node bifurcation. We are now interested in stationary distributions of the saddle-node equation distinct from $\delta_{\sqrt{a}}$ and $\delta_{-\sqrt{a}}$. Since the system presents two singular points, the results of proposition 5, based on integrability properties on $\mathbb{R}^+$ or $\mathbb{R}^-$ of the solutions of the Kolmogorov equation, do not directly apply. However, a similar approach allows to demonstrate the following:

Proposition 9. *Stationary distributions of the saddle-node equation* (14) *distinct from* $\delta_{\sqrt{a}}$ *and* $\delta_{-\sqrt{a}}$ *(when* $a > 0$*) enjoy the following classification:*

- *For* $\alpha = 1$ *and* $0 < a < \sigma^{-4}$, *the system has a stationary distribution, stable in the first approximation, charging only the interval* $(-\infty, -\sqrt{a})$ *that undergoes a P-bifurcation at* $\sigma^2\sqrt{a} = \frac{1}{2}$. *On* $(-\sqrt{a}, \infty)$, *any solution converges almost surely exponentially towards* $\sqrt{a}$ *(see Figure 9).*
- *For* $\alpha = 1$ *and* $a < 0$, *there exists a unique distribution charging the whole real line.*
- *For* $\alpha > 1$, *the system has no additional stationary solution.*

Proof. In the case $\alpha = 1$, the Kolmogorov equation has the solution:

$$p_0(x) \propto \left|x + \sqrt{a}\right|^{-2+\frac{1}{\sigma^2\sqrt{a}}} \left|x - \sqrt{a}\right|^{-2-\frac{1}{\sigma^2\sqrt{a}}}$$

For $a > 0$, this function is integrable at $\sqrt{a}$ if and only if $-2-\frac{1}{\sigma^2\sqrt{a}} > -1$ which is not possible. At $-\sqrt{a}$, this function is integrable when $-2 + \frac{1}{\sigma^2\sqrt{a}} > 1$ i.e. $\sigma^2\sqrt{a} < 1$ (i.e. when $-\sqrt{a}$ is unstable). Eventually, this function is integrable at infinity whatever the parameters, since it behaves like x^{-4}. Hence there exists a unique stationary distribution charging $(-\infty, -\sqrt{a})$. In both the interval $(-\sqrt{a}, \sqrt{a})$ and $(\sqrt{a}, \infty)$ the only stationary distribution is $\delta_{\sqrt{a}}$ which is stable. The additional stationary distribution on $(-\infty, -\sqrt{a})$ reads:

$$p_0(x) = Z\left(-x - \sqrt{a}\right)^{-2+\frac{1}{\sigma^2\sqrt{a}}} \left(-x + \sqrt{a}\right)^{-2-\frac{1}{\sigma^2\sqrt{a}}} =: Z\, q_0(x)$$

Denoting $\Phi = \frac{1}{\sigma^2\sqrt{a}}$, we give the following closed form primitives related:

$$\begin{cases} \displaystyle\int q_0(x)\,dx & = -(-\sqrt{a} - x)^{-1+\Phi}(\sqrt{a} - x)^{-1-\Phi}\dfrac{a(2\Phi^2 - 1) - 2\sqrt{a}\Phi x + x^2}{4a^{3/2}\Phi(\Phi^2 - 1)} \\ \displaystyle\int x\,q_0(x)\,dx & = (-\sqrt{a} - x)^{-1+\Phi}(\sqrt{a} - x)^{-1-\Phi}\dfrac{-2\sqrt{a}\Phi x + a + x^2}{4a(\Phi^2 - 1)} \\ \displaystyle\int x^2\,q_0(x)\,dx & = -(-\sqrt{a} - x)^{-1+\Phi}(\sqrt{a} - x)^{-1-\Phi}\dfrac{-2\sqrt{a}\Phi x + a + (2\Phi^2 - 1)x^2}{4\sqrt{a}\Phi(\Phi^2 - 1)} \end{cases}$$

Using these formulae, it is straightforward to obtain the normalization constant of p_0 on $(-\infty, -\sqrt{a})$:

$$Z = 4a^{3/2}\Phi(\Phi^2 - 1)$$

The expectation m is equal to:

$$m = Z\int_{-\infty}^{-\sqrt{a}} xq_0(x)\,dx = -\Phi\sqrt{a} = -\frac{1}{\sigma^2},$$

and the second moment:

$$s = Z\int_{-\infty}^{-\sqrt{a}} x^2 q_0(x)\,dx = (2\Phi^2 - 1)a$$

Eventually, let us analyze the shape of the obtained stationary distribution. The differential of the distribution reads:

$$p_0'(x) = p_0(x)\left(\frac{-2-\Phi}{x-\sqrt{a}} + \frac{-2+\Phi}{x+\sqrt{a}}\right).$$

This quantity is always negative for $\Phi < 2$, the density is decreasing and moreover in that case it is clear that the distribution diverges at $-\sqrt{a}$. For $\Phi > 2$, the distribution vanishes at zero, takes its maximum at $x^* = -\frac{\Phi\sqrt{a}}{2} = -\frac{1}{2\sigma^2}$ and goes back to zero.

Let us now address the stability in the first approximation for this distribution. The linearized equation reads:

$$dv_t = -2X_t v_t\,dt + 2\sigma X_t v_t dW_t$$

Integrating the linearized equation as previously done, we obtain that the Lyapunov exponent of the linearized equation reads:

$$l = -2\mathbb{E}[X_t] - 2\sigma^2\mathbb{E}[X_t^2] = \frac{2(-1+\sigma^4 a)}{\sigma^2}$$

and since we consider the case $\sigma^2\sqrt{a} < 1$, the Lyapunov exponent is strictly negative, implying almost sure exponential stability of zero for the linearized equation, hence stochastic linear stability of $p_0(x)$.

For $a < 0$, the solutions of the Kolmogorov equation are given by:

$$p_0(x) = \frac{K}{(x^2-a)^2}\exp\left(-2\frac{\arctan(x/\sqrt{-a})}{\sigma^2\sqrt{-a}}\right)$$

and the definition interval of possible stationary distributions is $\mathbb{R}$. In order for p_0 to define a probability distribution, we need this function to be integrable at $\pm\infty$, which is always the case. The obtained stationary distribution reaches its maximum again for $x = -\frac{1}{2\sigma^2}$.

For $\alpha > 1$, the solution of the Kolmogorov equation reads:

$$p(x) = \frac{1}{(a-x^2)^{2\alpha}}\exp\left(\frac{2}{\sigma^2}\int_{.}^{x}(-y^2+a)^{1-2\alpha}\,dy\right)$$

At $-\sqrt{a}$, the exponential term diverges since it behaves as $(x+\sqrt{a})^{1-2\alpha}$, hence the solution is never integrable at zero. It is easy at this point to show that there is no integrable solution to the Kolmogorov equation. □

Remark. *For $\alpha < 1$ one could be interested in the existence of quasi-stationary solutions. An additional difficulty arises from the fact that the solutions can be attracted in finite time towards different solutions: $\pm\sqrt{a}$ whatever $\alpha < 1$, and $-\infty$ for $\alpha < \frac{3}{4}$. Moreover, in the saddle-node bifurcation case as studied here, the diffusion coefficient is exactly equal to a power function. Therefore, for $\alpha < 1$, blow up in finite time as well as absorption at $\pm\sqrt{a}$ can occur are not a consequence of the results of [15], are not in the scope of the present manuscript and require new mathematical developments that will be addressed elsewhere.*



Figure 9: Behaviors of the stochastic saddle-node system with $\alpha = 1$. The bifurcation diagram as a function of a is at the center of the figure, and displays the singular points in blue lines, their stability marked by the type of curve: plain: stable, dashed: unstable. Stationary distributions are encoded in color in the interval $a \leq \sigma^{-4}$. Two typical stationary distributions are plotted in Figs. (e) and (f): one for $a \leq 1/4\sigma^4$ where the pdf has a peak at $x = -\sigma^{-2}/2$, and for $a > \sigma^{-4}/4$ where the distribution diverges at zero. The P-bifurcation between these behaviors is noted P. (a) corresponds to a typical trajectory for $a = 0.4 < \sigma^{-4}/4$ with $\sigma = 0.8$. (b),(c) and (d) correspond to $a > \sigma^{-4}$ and different initial conditions: (c): frequency of the trajectories converging towards $+\sqrt{a}$ as a function of the initial condition in $[-\sqrt{a}, \sqrt{a}]$, (b) and (d) are typical trajectories for initial condition in $(\sqrt{(a)}, \infty)$ or $(-\infty, -\sqrt{a})$ respectively ($a = 4$ and $\sigma = 0.8$).

The above result characterize the dynamics relatively exhaustively for $\alpha = 1$, except for $a > \sigma^{-4}$ and $x_0 \in (-\sqrt{a}, \sqrt{a})$. In that case, both singular points are stable, and can be reached by the solution. Therefore for such initial conditions, there is a competition between the possible attractors, and the relative stability of one singular point compared to the other will determine the probability, given an initial condition x_0, to converge towards this equilibrium. This quantity is plotted in Figure 9(c). We observe, as one can expect, that the attractivity of $\sqrt{a}$ is stronger than that of $-\sqrt{a}$: the probability of converging towards $\sqrt{a}$ is never less than 0.4 even for initial conditions very close from $-\sqrt{a}$, and goes relatively fast towards 1 as the initial condition is chosen closer from $\sqrt{a}$.

Let us now discuss the universality of this bifurcation. We consider a general one-dimensional diffusion equation:

$$dx_t = f(x_t, \lambda)\,dt + g(x_t)dW_t$$

with f a smooth function such that $f(0,0) = \frac{\partial f}{\partial x}(0,0) = 0$ and $\frac{\partial f}{\partial \lambda}(0,0) \neq 0$. Following the reduction to a normal form of Kuznetsov [2], we use a Taylor expansion of f:

$$f(x,\lambda) = f_0(\lambda) + f_1(\lambda)\,x + f_2(\lambda)\,x^2 + h(x)$$

where $h(x)/x^3$ is bounded at $x = 0$. We perform a shift of coordinates by defining a new variable $\xi = x + \delta$ and obtain:

$$\begin{aligned} d\xi_t = \Big([f_0(\lambda) - f_1(\lambda)\delta] + f_2(\lambda)\delta^2 + O(\delta^3)]\\ + [f_1(\lambda) - 2f_2(\lambda)\delta + O(\delta^2)]\xi\\ + [f_2(\lambda) + O(\delta)]\xi^2 + O(\xi^3)\Big)\,dt + g(\xi - \delta)dW_t \end{aligned}$$

Using the inverse function theorem under the assumption that $\frac{\partial f}{\partial \lambda}(0,0) \neq 0$, we deduce that there exists a smooth function $\delta(\lambda) = \frac{f_1'(0)}{2f_2(0)}\lambda + O(\lambda^2)$ such that the term $[f_1(\lambda) - 2f_2(\lambda)\delta + O(\delta^2)]$ vanishes for all λ in a neighborhood of 0. This $\delta(\lambda)$ identified, we can apply the inverse function theorem to find locally an inverse to the constant term of the Taylor development seen as a function of λ. This inverse function, $\lambda(\mu)$, yields the system to the SDE:

$$d\xi_t = (\mu + b(\mu)\xi_t^2 + O(\xi^3))\,dt + g(\xi - \delta(\lambda(\mu)))dW_t$$

with $b(0) = f_2(0) \neq 0$. The final reduction involves a scaling of the variable: $z_t = |b(\mu)|\xi_t$, yielding the SDE:

$$dz_t = \Big(\beta + \varepsilon z_t^2 + O(z_t^3)\Big)\,dt + g\Big(\frac{z_t}{|b(\mu)|} - \delta(\lambda(\mu))\Big)dW_t$$

where ε has the sign of $S^{(3)}(0)$. The stability and bifurcations of the whole system is hence reduced to the local analysis of the behavior of g close of the equilibria.

Example 3. *We illustrate this method using a specific instance of a one-population Wilson and Cowan system with constant external input $I \in \mathbb{R}$:*

$$dv_t = \Big(-v_t + S(gv_t) + I\Big)\,dt + \sigma|-v_t + S(gv_t) + I|^{\alpha}dW_t$$

and choose as usual $S(x) = \tanh(g\,x)$. The whole dynamics is hence governed by a single function $f(x) = -x + S(g\,x) + I$ and the deterministic system undergoes two saddle-node bifurcation when

$g > 1$, located at $v = \pm\frac{\sqrt{g-1}}{g}$ for an input equal to (respectively): $I = \pm\frac{\sqrt{g-1}}{g} \mp S(\sqrt{g-1})$. Let us focus for instance on the bifurcation arising at $\frac{\sqrt{g-1}}{g}$. Around this point, a Taylor expansion yields, denoting $x_t = v_t - \frac{\sqrt{g-1}}{g}$:

$$dx_t = \left(\left(I - \frac{\sqrt{g-1}}{g} + S(\sqrt{g-1})\right) - \sqrt{g-1}\, x_t^2 + h(x_t)\right) dt \\ + \sigma \left|\left(I - \frac{\sqrt{g-1}}{g} + S(\sqrt{g-1})\right) - \sqrt{g-1} x_t^2 + h(x_t)\right|^{\alpha} dW_t$$

where $h(x) = O(x^3)$ at zero. Eventually, denoting $\xi_t = \sqrt{g-1}\, x_t$ we obtain:

$$d\xi_t = (\beta - \xi_t^2 + \Psi(\xi_t))\, dt + |\beta - \xi_t^2 + \Psi(\xi_t)|^{\alpha} dW_t$$

with $\beta = \sqrt{g-1}(I + S(\sqrt{g-1}) - \frac{g-1}{g}$ and $\Psi(x) = \sqrt{g-1}\, h(\frac{x}{\sqrt{g-1}})$. The system is hence locally reduced to the saddle-node equation, and the above analysis characterizes the dynamics of the system. Let us for instance consider I as our parameter. As I crosses $I^* = \frac{\sqrt{g-1}}{g} - S(\sqrt{g-1})$, a stochastic saddle-node bifurcation occurs. For $I > I^*$ the system has two equilibria $x_l < \sqrt{g-1} < x_r$ and an additional singular point x^* strictly smaller than x_l. It is easy to show that this singular point is always stable in probability. Regarding x_l and x_r, we have:

- *For $\alpha < 1$, both x_l and x_r are stable and the first hitting time of one of the equilibria is finite almost surely.*
- *For $\alpha = 1$, x_r is always stable in probability, and x_l is stable for $\beta > \sigma^{-4}$, and unstable otherwise.*
- *For $\alpha > 1$, x_r is stable in probability and x_l unstable.*

We illustrate this variety of behaviors in Figure 10.

4. Discussion

In this manuscript we analyzed the interplay between noise and nonlinearities in the setting of SDEs with singular points, and showed that the present analysis could account for the dynamics of a number of low-dimensional stochastic systems driven by a unidimensional Brownian motion.

Even though these limitations restrict the variety of behaviors possible, we observed and quantified several effects of noise, such as for instance the that it can prevent explosion of the solutions, and that it interferes with the stability of singular points. Moreover, as soon as the dimension of the vector field is increased (e.g. Hopf bifurcation), stochastic perturbations can have more complex effects in which the precise structure of the noise interacts subtly with the dynamics. In particular we identified a new bifurcation corresponding to a change in the orientation rotation of the stochastic cycles.

An important question related to this manuscript is the possibility to reduce the dynamics of general SDEs to low-dimensional systems, analogous to the center manifold reduction of



(a) $\alpha = 0.8, \sigma = 2$

(b) $\alpha = 0.8$, $\sigma = 2$ (other sample path)

(c) $\alpha = 1, \sigma = 2$



(d) $\alpha = 1$, $\sigma = 2$ (other sample path)

(e) $\alpha = 1, \sigma = 0.2$

(f) $\alpha = 1.5, \sigma = 0.2$

Figure 10: Dynamics of the Wilson and Cowan System (v_t as a function of time) for different values of α in the saddle-node case with $g = 3$, $I = 0$, and initial condition $\frac{\sqrt{g-1}}{g}$. In that case, $x_r = 0$ and x_l has no closed-form expression. We illustrate the stabilization of x_l in the case $\alpha \leq 1$, which holds only for large noise values when $\alpha = 1$: (a) and (b): $\alpha < 1$, x_l and x_r are stable in probability and the first hitting time to one of the equilibria is a.s. finite. (c) $\alpha = 1$, σ large enough so that both equilibria are stable, (d): for σ small, only x_l is stable. (d): for $\alpha > 1$, only x_l is stable. The initial condition is taken in the interval (x_r, x_l), and the stability of x^* is not addressed.

deterministic dynamical systems. This is highly dependent on the type of diffusion coefficients considered, and the dimension of the Brownian motion involved in the dynamics. Indeed, in the stochastic case, when performing nonlinear changes of variables, additional terms appear in the drift function, that depend on both the drift and diffusion coefficients. This analysis is part of the perspectives of this work. Another perspective is the further analysis of the quasi-stationary distributions of such systems. Indeed, the present analysis demonstrated that quasi-stationary distributions existed as long as $\alpha < \frac{3}{4}$. The question hence arises whether this finding is a limitation of the framework we used, or if actually disappears and we are left with no stationary behavior for the trajectories that did not reach zero. This would be the first kind of process discovered, up to our knowledge, of diffusion absorbed in finite time, with no quasi-stationary distribution. We believe that in the case $\lambda > 0$, the quasi-stationary distributions progressively turn into stationary distributions as the exponent α increases to 1. This would correspond to a convergence towards zero of the eigenvalue of the Kolmogorov operator related to the quasi-stationary distribution. But for $\lambda < 0$ and $\alpha > 1$, there is no stationary distribution for the process, and hence these quasi-stationary distribution do disappear. And numerical simulations provided in Figure 2 tend to confirm that indeed, the quasi-stationary distribution disappear in favor of a progressive convergence towards zero of all trajectories conditioned on not hitting zero. The analysis of this phenomenon is another perspective for future work.

References

[1] J. Guckenheimer, P. J. Holmes, Nonlinear Oscillations, Dynamical Systems and Bifurcations of Vector Fields, volume 42 of *Applied mathematical sciences*, Springer, 1983.

[2] Y. A. Kuznetsov, Elements of Applied Bifurcation Theory, Applied Mathematical Sciences, Springer, 2nd edition, 1998.

[3] L. Arnold, H. Crauel, V. Wihstutz, Stabilization of linear systems by noise, SIAM Journal on Control and Optimization 21 (1983) 451.

[4] J. Pham, K. Pakdaman, J. Vibert, Noise-induced coherent oscillations in randomly connected neural networks, Physical Review E 58 (1998) 3610.

[5] C. Peskin, G. Odell, G. Oster, Cellular motions and thermal fluctuations: the brownian ratchet., Biophysical Journal 65 (1993) 316.

[6] C. Doering, W. Horsthemke, J. Riordan, Nonequilibrium fluctuation-induced transport, Physical review letters 72 (1994) 2984–2987.

[7] L. Gammaitoni, P. Hänggi, P. Jung, F. Marchesoni, Stochastic resonance, Reviews of Modern Physics 70 (1998) 223.

[8] A. Pikovsky, J. Kurths, Coherence resonance in a noise-driven excitable system, Physical Review Letters 78 (1997) 775–778.

[9] L. Arnold, Random dynamical systems, Springer Verlag, 1998.

[10] H. Crauel, P. Imkeller, M. Steinkamp, Bifurcations of one-dimensional stochastic differential equations, Stochastic dynamics (1999) 27–47.

[11] S. Tanabe, K. Pakdaman, Dynamics of moments of fitzhugh-nagumo neuronal models and stochastic bifurcations, Physical Review E 63 (2001) 031911.

[12] J. Touboul, G. Hermann, O. Faugeras, Noise-induced behaviors in neural mean field dynamics, SIAM J. on Dynamical Systems 11 (2011).

[13] N. Berglund, B. Gentz, Geometric singular perturbation theory for stochastic differential equations, Journal of Differential Equations 191 (2003) 1–54.

[14] G. Pavliotis, A. Stuart, Multiscale methods: averaging and homogenization, Springer Verlag, 2008.

[15] P. Cattiaux, P. Collet, A. Lambert, S. Martinez, S. Méléard, J. San Martín, Quasi-stationary distributions and diffusion models in population dynamics, The Annals of Probability 37 (2009) 1926–1969.

[16] K. Pakdaman, M. Thieullen, G. Wainrib, Fluid limit theorems for stochastic hybrid systems with application to neuron models, Advances in Applied Probability 42 (2010) 761–794.

[17] R. Has'minskii, Stochastic stability of differential equations, Kluwer Academic Pub, 1980.

[18] X. Mao, Stochastic Differential Equations and Applications, Horwood publishing, 2008.

[19] H. Wilson, J. Cowan, Excitatory and inhibitory interactions in localized populations of model neurons, Biophys. J. 12 (1972) 1–24.
[20] D. Revuz, M. Yor, Continuous martingales and Brownian motion, Springer Verlag, 1999.
[21] I. Karatzas, S. Shreve, Brownian motion and stochatic calculus, Springer, 1987.
[22] M. Freidlin, A. Wentzell, Random perturbations of dynamical systems, Springer Verlag, 1998.